

# Sufficient conditions for forward invariance and contractivity in hybrid inclusions using barrier functions

Mohamed Maghenem<sup>a</sup> and Ricardo G. Sanfelice<sup>a</sup>

<sup>a</sup>*Department of Electrical and Computer Engineering, University of California, Santa Cruz.*

---

## Abstract

This paper studies set invariance and contractivity in hybrid systems modeled by hybrid inclusions using barrier functions. After introducing the notion of barrier function, we investigate sufficient conditions to guarantee different forward invariance and contractivity notions of a closed set for hybrid systems with nonuniqueness of solutions and solutions terminating prematurely. We consider forward (pre-)invariance of sets, which guarantees that the maximal solutions starting from the set stay in it, and (pre-)contractivity, which further requires that the solutions starting from the boundary of the set evolve immediately (continuously or discretely) towards its interior. Our conditions for forward invariance and contractivity are infinitesimal and in terms of the proposed barrier functions. Examples illustrate the results.

*Key words:* Forward invariance, contractivity, barrier functions, hybrid dynamical systems.

---

## 1 Introduction

Forward invariance of sets for dynamical systems is a property that requires the solutions starting in the considered set to remain in it along their entire domain of definition. The main challenge when studying forward invariance consists of providing sufficient conditions while avoiding explicit computation of the system solutions.

The study of set invariance for dynamical systems is a key step towards analyzing their stability and safety properties. Indeed, forward invariance has a close relationship to safety, which is a property that requires the system solutions starting from a given set of initial conditions to remain in a desired safe region [1]. Safety, also named conditional invariance in [2], is equivalent to forward invariance of a set, known as *inductive invariant* in [3], which contains the set of initial conditions and not intersecting with the unsafe region [4]. Furthermore,

the study of set invariance can be a key step to conclude stability properties for the system via relaxed Lyapunov conditions, including the well-known *invariance principle* [5] and *Matrosov Theorem* [6]. In addition to safety, the study of set invariance has also been extended in order to guarantee some closely related notions such as *quasi-invariance*, *conditional quasi-invariance* [7], and *contractivity* [8]. The contractivity property is a strong form of forward invariance. Indeed, a contractive set is forward invariant, and whenever a solution starts from its boundary, it immediately leaves the boundary and evolves towards its interior. Contractivity is also named *strict invariance* in [9]. Analyzing contractivity, which is a strong form of forward invariance, in the context of barrier functions is useful. Indeed, contractivity is closely related to invariance and it has been studied in the literature, for example, when computing *set-induced Lyapunov functions* [10–12]. Hence, our goal is to formalize the differences between the two notions and how to certify each of them using barrier functions.

### 1.1 Background

The interest in the study and characterization of forward invariance, while avoiding the computation of the system's solutions, dates back to the seminal work of Nagumo in [13]. In this reference, conditions involving the contingent cone and the system's dynamics on the boundary of a closed set are shown to be necessary and sufficient to conclude, from each point in the set, the ex-

---

*Email addresses:* mmaghene@ucsc.edu (Mohamed Maghenem), ricardo@ucsc.edu (Ricardo G. Sanfelice).

This research has been partially supported by the National Science Foundation under Grant no. ECS-1710621 and Grant no. CNS-1544396, by the Air Force Office of Scientific Research under Grant no. FA9550-16-1-0015, Grant no. FA9550-19-1-0053, and Grant no. FA9550-19-1-0169, and by CITRIS and the Banatao Institute at the University of California.

istence of at least one solution that remains in the set. This last property defines what is known as *weak forward invariance* [14], which is equivalent to forward invariance when the system’s solutions are unique. Extensions of this result, using a similar type of cone conditions, are presented in [9] to conclude weak forward invariance<sup>1</sup> for differential inclusions, in [15] for impulse differential inclusions, and in [16] for hybrid inclusions. For systems with continuous-time dynamics, when all the solutions starting from a closed set are required to remain in it, as stressed in [9], the invariance conditions concern the system’s dynamics outside the set rather than on its boundary. As a consequence, the external contingent cone is introduced and used in [9].

The relatively stronger form of invariance called contractivity is characterized in [8] in terms of the *Minkowski* functional, both for differential and difference equations, for the particular case of convex and compact sets. For general closed sets and for systems modeled as differential inclusions, sufficient conditions are proposed in [9] using, roughly speaking, the interior of the contingent cone and the system’s dynamics at the boundary of the considered closed set. In general, the computation of the tangent cones is not a trivial task. However, when the considered set is defined as the zero-sublevel of a vector function, named *barrier function candidate*, it is possible, under appropriate assumptions, to formulate invariance and contractivity conditions using only the barrier function candidate and the system’s dynamics.

The latter approach has been adopted in [17] and [18, 19] for differential equations and inclusions, respectively, and in [20] for hybrid automata. In [17], another type of barrier function candidate is also considered. Such a barrier function candidate is positive and locally bounded on the interior of the considered set, and approaches infinity as its argument converges to the boundary of the set. When using the latter notion of barrier functions, solutions starting from the interior of the set to render invariant are not allowed to reach its boundary. In [18], sufficient conditions for invariance in terms of nonsmooth but locally Lipschitz barrier functions are considered with application to safe navigation for networks of vehicles in the presence of obstacles. See also [21, 22] for more applications. Finally, in [20], methods to synthesize barrier functions are investigated. The latter work is extended in [23, 24] by relaxing the conditions constraining the continuous-time evolution of the hybrid automata.

## 1.2 Motivation

A hybrid inclusion is defined as a differential inclusion with a constraint, which models the *flow* or continuous evolution of the system, and a difference inclusion with a constraint, modeling the *jumps*, or discrete events.

<sup>1</sup> Weak forward invariance is named *viability* in [9].

In particular, handling nonuniqueness of solutions in hybrid inclusions and solutions terminating prematurely lead to particular forms of forward invariance and contractivity properties that we call forward pre-invariance and pre-contractivity, respectively, where the prefix “pre” indicates that some solutions may have a bounded (hybrid) time domain. The aforementioned notions have not been covered in the literature using barrier functions. Furthermore, having sufficient conditions for forward invariance in terms of barrier functions is useful, especially when control inputs are used to force such conditions [1, 17], or when the invariance, or the contractivity, task is to be combined with a stabilization task to be achieved inside a safety set [25]. Furthermore, in many applications, it is often the case that the closed set to be rendered forward invariant or contractive corresponds to the region where multiple scalar functions are nonpositive simultaneously. In such a case, it is typically difficult to find a single scalar function that defines the set of interest and, at the same time, is sufficiently smooth. This fact motivates the development of sufficient conditions guaranteeing forward invariance and contractivity when multiple scalar candidates define the considered set.

## 1.3 Contributions

In this paper, we introduce barrier functions and tools to certify forward invariance and contractivity in hybrid systems modeled as hybrid inclusions. We define a barrier function candidate as a vector function of the state variables. Sufficient conditions in terms of infinitesimal inequalities – namely, without using information about solutions – are proposed to guarantee that the set of points on which all the components of the barrier function candidate are nonpositive is forward invariant or contractive. More precisely, under mild conditions on the data defining the hybrid inclusion, we present conditions such that a barrier function candidate guarantees forward pre-invariance. The proposed conditions can be decomposed into flow and jump conditions that restrict the continuous and the discrete evolution of the hybrid system, respectively.

In Section 3, sufficient conditions for forward invariance that apply when the barrier function candidate is continuously differentiable and only locally Lipschitz are presented; see, respectively, Theorems 1 and 4. Furthermore, the flow conditions therein need to be satisfied on a neighborhood outside the considered set. The flow conditions in Theorems 1 and 4 are relaxed in Proposition 1 and Remark 13 using uniqueness functions. This relaxation is original to the best of our knowledge. On the other hand, under a regularity condition on the gradient of the barrier function candidate, known as *transversality condition*, which is typically assumed in the literature (see, e.g., [9, 26]), plus some extra regularity conditions on the flow dynamics, in Theorem 2, the flow condition

needs to hold only on a smaller region of the state space which is the boundary of the considered set.

In Section 4, we analyze contractivity properties for hybrid systems. After a brief overview of the case of sets that are compact and convex, which itself extends to hybrid inclusions the results in [8] for differential and difference equations, we introduce a notion of contractivity for general closed sets. Furthermore, it extends to hybrid inclusions what is proposed in [9] for differential inclusions only. The proposed notion essentially requires the system's solutions to evolve from points on the boundary of  $K$  towards its interior via a flow or a jump. Sufficient conditions for contractivity in terms of the barrier function candidates defining the set are established when the latter candidate is either continuously differentiable or only locally Lipschitz, see Theorems 5 and 6.

The results in this paper extend what was proposed in [17, 18, 20, 23, 24] to the more general context of hybrid systems modeled by hybrid inclusions. That is, hybrid inclusions offer many technical challenges that have not been handled in the existing literature. Those challenges are mainly due to the fact that the continuous-time evolution of the hybrid inclusion is not necessarily defined on an open set. Moreover, the considered set is defined as the zero-sublevel set of a barrier function candidate restricted to the set where the dynamics are defined; namely, the union of the flow and the jump sets. Since the latter sets can be closed, elements around the intersection between the two boundaries; namely, the zero-level set of the barrier candidate and the boundary of the flow set, needs a particular treatment. Our sufficient conditions using barrier functions are alternatives to those proposed in [16] and [9] using tangent cone-based conditions. Indeed, our conditions exploit the fact that the set is the intersection of zero-sublevel sets of scalar functions; hence, the obtained conditions avoid as much as possible the computation of tangent cones. The latter task is known to be numerically expensive in some cases. It is also to be noted that some of our results build upon the well-known cone-based conditions in [9] and [14].

To the best of our knowledge, this is the first time in the literature where the concept of barrier functions is used for hybrid inclusions to analyze different set-invariance properties. Preliminary versions of this work are in the conference papers [27, 28]. However, only scalar barrier functions are considered in [29], and many proofs, explanations, and examples are omitted in both submissions. Furthermore, compared with [27, 28], new results are proposed in Theorem 2, Theorem 3, Theorem 4, Proposition 1, and Theorem 6.

The remainder of the paper is organized as follows. Preliminaries and basic conditions are presented in Section 2. Sufficient conditions of forward pre-invariance and invariance using barrier functions are in Section 3. Sufficient conditions for pre-contractivity and contractivity

using barrier functions are in Section 4, respectively. Examples are included at each step in order to illustrate the proposed statements. More technical details can be found in [30].

**Notation.** Let  $\mathbb{R}_{\geq 0} := [0, \infty)$ ,  $\mathbb{N} := \{0, 1, \dots\}$ , and  $\mathbb{N}^* := \{1, 2, \dots, \infty\}$ . For  $x, y \in \mathbb{R}^n$  and a nonempty set  $K \subset \mathbb{R}^n$ ,  $x^\top$  denotes the transpose of  $x$ ,  $|x|$  the norm of  $x$ ,  $|x|_K := \inf_{y \in K} |x - y|$  defines the distance between  $x$  and the set  $K$ ,  $\langle x, y \rangle = x^\top y$  denotes the inner product between  $x$  and  $y$ , and  $\langle x, K \rangle = x^\top K := \{x^\top z : z \in K\}$ . The inequalities  $x \leq 0$  and  $x < 0$  mean that  $x_i \leq 0$  and, respectively,  $x_i < 0$  for all  $i \in \{1, 2, \dots, n\}$ . The opposites, namely,  $x \not\leq 0$  and  $x \not< 0$  mean that there exists  $i \in \{1, 2, \dots, n\}$  such that  $x_i > 0$  and, respectively,  $x_i \geq 0$ . For a set  $K \subset \mathbb{R}^n$ , we use  $\text{int}(K)$  to denote its interior,  $\partial K$  to denote its boundary,  $\text{cl}(K)$  to denote its closure, and  $U(K)$  to denote an open neighborhood around  $K$ . For a set  $O \subset \mathbb{R}^n$ ,  $K \setminus O$  denotes the subset of elements of  $K$  that are not in  $O$ . By  $\mathbb{B}$ , we denote the closed unit ball in  $\mathbb{R}^n$  centered at the origin. For a function  $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$  and a set  $D \subset \mathbb{R}^m$ ,  $f(D) = \{f(x) : x \in D\}$ . For a continuously differentiable function  $B : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $\nabla B(x)$  denotes the gradient of the function  $B$  evaluated at  $x$ . By  $\mathcal{C}^1$ , we denote the set of continuously differentiable functions. Finally,  $F : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$  denotes a set-valued map associating each element  $x \in \mathbb{R}^n$  into a subset  $F(x) \subset \mathbb{R}^n$ .

## 2 Preliminaries and Basic Conditions

### 2.1 Hybrid Inclusions

We consider hybrid systems modeled by

$$\mathcal{H} : \begin{cases} x \in C & \dot{x} \in F(x) \\ x \in D & x^+ \in G(x), \end{cases} \quad (1)$$

with the state variable  $x \in \mathbb{R}^n$ , the flow set  $C \subset \mathbb{R}^n$ , the jump set  $D \subset \mathbb{R}^n$ , the flow and the jump set-valued maps, respectively,  $F : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ ,  $G : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ . A solution  $x$  to  $\mathcal{H}$  is defined on a hybrid time domain denoted  $\text{dom } x \subset \mathbb{R}_{\geq 0} \times \mathbb{N}$ . The solution  $x$  is parametrized by the ordinary time variable  $t \in \mathbb{R}_{\geq 0}$  and the discrete jump variable  $j \in \mathbb{N}$ . Its domain of definition  $\text{dom } x$  is such that for each  $(T, J) \in \text{dom } x$ ,  $\text{dom } x \cap ([0, T] \times \{0, 1, \dots, J\}) = \cup_{j=0}^J ([t_j, t_{j+1}], j)$  for a sequence  $\{t_j\}_{j=0}^{J+1}$ , such that  $t_{j+1} \geq t_j$  for each  $j \in \{0, 1, \dots, J\}$  and  $t_0 = 0$ ; see [31].

A solution  $x$  to  $\mathcal{H}$ , as defined in [30, Definition 12], starting from  $x_o$  is said to be complete if it is defined on an unbounded hybrid time domain; that is, the set  $\text{dom } x$  is unbounded. Furthermore, it is said to be maximal if there is no solution  $y$  to  $\mathcal{H}$  such that  $x(t, j) = y(t, j)$  for all  $(t, j) \in \text{dom } x$  with  $\text{dom } x$  a proper subset of  $\text{dom } y$ .

Finally, it is said to be nontrivial if  $\text{dom } x$  includes at least two points. To simplify notation, we use  $x$  to denote the state and also a solution. We write “solution  $x$ ” to make this distinction when needed.

## 2.2 Anatomy of Sets

Different types of cones have been used in the study of differential inclusions. In the following, for a set  $K \subset \mathbb{R}^n$ , we recall the ones we use in this paper, see [9] for more details. The *contingent* cone of  $K$  at  $x$  is given by

$$T_K(x) := \left\{ v \in \mathbb{R}^n : \liminf_{h \rightarrow 0^+} \frac{|x + hv|_K}{h} = 0 \right\}. \quad (2)$$

We also recall the equivalence [32, Page 122]

$$v \in T_K(x) \Leftrightarrow \exists \{h_i\}_{i \in \mathbb{N}} \rightarrow 0^+ \text{ and } \{v_i\}_{i \in \mathbb{N}} \rightarrow v : \\ x + h_i v_i \in K \quad \forall i \in \mathbb{N}. \quad (3)$$

The *external contingent* cone of  $K$  at  $x$  is given by

$$E_K(x) := \left\{ v \in \mathbb{R}^n : \liminf_{h \rightarrow 0^+} \frac{|x + hv|_K - |x|_K}{h} \leq 0 \right\}. \quad (4)$$

The *Dubovitsky-Miliutin* cone of  $K$  at  $x$  is given by

$$D_K(x) := \{v \in \mathbb{R}^n : \exists \epsilon > 0 : \\ x + \delta(v + w) \in K \quad \forall \delta \in (0, \epsilon], \forall w \in \epsilon \mathbb{B}\}. \quad (5)$$

## 2.3 Basic Assumptions

The proposed results on forward invariance and contractivity of a set  $K \subset C \cup D$  for  $\mathcal{H}$  are obtained under the following standing assumption.

**Standing assumption.** The data of the hybrid inclusion  $\mathcal{H} = (C, F, G, D)$  is such that the flow map  $F$  is outer semicontinuous and locally bounded with nonempty and convex images on  $C$ , and  $G(x)$  is nonempty for all  $x \in D$ . Furthermore, the set  $K$  is closed. •

We notice that, in addition to these standing assumptions, the hybrid basic conditions in [33, Chapter 6], which are not imposed here, also require the sets  $C$  and  $D$  to be closed and the jump map  $G$  to be locally bounded.

Before going further, consider the hybrid inclusion  $\mathcal{H} = (C, F, D, G)$  and a closed set  $K \subset C \cup D$ . Starting from  $x_o \in K$ , if a solution  $x$  leaves the set  $K$ , then it has to be under one of the two following scenarios:

(Sc1) The solution  $x$  leaves the set  $K$  after a jump. It implies the existence of  $(t, j) \in \text{dom } x$  such that  $x(t, j) \in K \cap D$  and  $(t, j+1) \in \text{dom } x$  with  $x(t, j+1) \notin K$  and  $x(t, j+1) \in G(x(t, j))$ .

(Sc2) The solution  $x$  leaves the set  $K$  by flowing. It implies the existence of  $t'_2 > t'_1 \geq 0$  and  $j' \in \mathbb{N}$  such that  $([t'_1, t'_2] \times \{j'\}) \subset \text{dom } x$  and  $x((t'_1, t'_2), j') \subset (U(\partial K) \setminus K) \cap C$ , with  $x(t'_1, j') \in \partial K$  and  $x(t'_2, j') \notin K$ .

In fact, when the set  $K$  is closed, under (Sc2),  $x(t'_1, j') \in \partial K \cap K$  and since the solution leaves the set  $K$ , under [30, Definition 12],  $x((t'_1, t'_2), j')$  is a subset of  $C \setminus K$  for some  $t'_2 > t'_1$  sufficiently close to  $t'_1$ .

## 3 Sufficient Conditions for Forward Pre-Invariance and Invariance Using Barrier Functions

Given a hybrid system  $\mathcal{H} = (C, F, D, G)$ , for a set  $K \subset C \cup D$ , following [16] and [33, Definition 6.25], we introduce the two following forward invariance notions.

**Definition 1 (Forward pre-invariance)** *The set  $K$  is said to be forward pre-invariant for  $\mathcal{H}$  if, for each  $x_o \in K$  and each maximal solution  $x$  starting from  $x_o$ ,  $x(t, j) \in K$  for all  $(t, j) \in \text{dom } x$ .*

**Definition 2 (Forward invariance)** *The set  $K$  is said to be forward invariant for  $\mathcal{H}$  if it is forward pre-invariant and for each  $x_o \in K$ , each maximal solution  $x$  starting from  $x_o$  is complete.*

Furthermore, we assume that the set  $K$  is defined as points in  $C \cup D$  at which multiple scalar functions are simultaneously nonpositive. These scalar functions form a barrier function candidate defining the set  $K$ .

**Definition 3** *A function  $B : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is said to be a barrier function candidate defining the set  $K$  if<sup>2</sup>*

$$K = \{x \in C \cup D : B(x) \leq 0\}, \quad (6)$$

where  $B(x) := [B_1(x) \ B_2(x) \ \dots \ B_m(x)]^\top$ .

If  $B$  is continuous, the set  $K$  is closed relative to  $C \cup D$ . If, in addition,  $C \cup D$  is closed, then  $K$  is automatically closed.

We introduce the following sets that we use in some statements and proofs. For a set  $K$  given as in (6), we define

$$K_e := \{x \in \mathbb{R}^n : B(x) \leq 0\}, \quad (7)$$

and, for each  $i \in \{1, 2, \dots, m\}$ ,

$$K_{ei} := \{x \in \mathbb{R}^n : B_i(x) \leq 0\}, \quad (8)$$

$$M_i := \{x \in \partial K : B_i(x) = 0\}. \quad (9)$$

<sup>2</sup>  $B(x) \leq 0$  means that  $B_i(x) \leq 0$  for all  $i \in \{1, 2, \dots, m\}$ .

It is useful to notice that  $K_e = \cap_{i=1}^m K_{ei}$ ,  $K = K_e \cap (C \cup D)$ , and that  $\partial K = \cup_{i=1}^m M_i \cup (\partial K \cap \partial(C \cup D))$ . Note that in general  $M_i \neq \partial K_{ei}$ .

**Remark 1** In existing literature, motivated by barrier methods for optimization [34], barrier function candidates<sup>3</sup> are introduced as scalar functions that are positive and locally bounded on  $\text{int}(K)$ , and approach infinity as their argument converges to  $\partial K$  [36]. The difference between the barrier function therein and the one in Definition 3 is that, the one in [36] guarantees that the solutions starting from  $\text{int}(K)$  cannot reach the boundary  $\partial K$ , which in turn renders  $\text{int}(K)$  invariant (in the appropriate sense), see [17] for a more detailed comparison.

**Remark 2** In the case of hybrid systems modeled as hybrid automata, the concept of barrier functions is used to conclude forward invariance, or safety in general, in [20, 23, 24]. According to these references, for a hybrid automata with  $m$  operating modes, a closed set  $K_q \subset \mathbb{R}^n$ ,  $q \in \{1, 2, \dots, Q\}$ , is associated to each mode (typically determined by the logic variable  $q$ ) and must be forward invariant for the state variable (typically denoted  $\zeta$ ) only during the corresponding  $q$  mode. The sets  $K_q$  can be different for each mode. Furthermore, since each mode is governed by a differential equation and the state variable  $\zeta$  is allowed to jump only when the mode switches, a barrier function candidate  $B_q$  is associated with each mode and defines the corresponding set  $K_q$  as  $K$  in Definition 3. Our approach covers such construction. Indeed, if we model a hybrid automata as a hybrid inclusion  $\mathcal{H}$  while incorporating the mode as a new discrete state variable  $q \in \{1, 2, \dots, Q\}$ , then, in the augmented space  $\mathbb{R}^n \times \{1, 2, \dots, Q\}$ , we can define the set  $K := \cup_{q=1}^Q (K_q \times \{q\})$  and the candidate  $B(\zeta, q) := B_q(\zeta)$ . It is easy to see that the latter scalar candidate  $B$  defines the set  $K$  according to Definition 3.

**Remark 3** In our approach, we do not restrict the barrier function candidate to be a scalar function. In general, one is interested in considering a forward invariant set  $K$  that is given by multiple inequality constraints being satisfied simultaneously. Also, we notice that it is always possible from (6) to construct a scalar barrier function candidate that defines the closed set  $K$  according to Definition 3 as

$$\bar{B}(x) := \max_{i \in \{1, 2, \dots, m\}} B_i(x). \quad (10)$$

However, by doing so, if the vector function  $B$  is  $\mathcal{C}^1$ , the resulting barrier function candidate  $\bar{B}$  is not guaranteed to be  $\mathcal{C}^1$  and it can be only continuous. Indeed, at points  $x$  where multiple  $B_i$ 's are equal, if their gradients are not identical, then  $\bar{B}$  is not differentiable at those elements.

<sup>3</sup> Barrier functions are also called *potential functions* in [35].

### 3.1 Pre-Invariance Under Standing Assumptions

The results we present in this section are extensions to what has been proposed in [20, 23, 24] for general hybrid inclusions while handling the possible noncomplete solutions and using multiple barrier functions instead of only a scalar one. For general differential inclusions (the continuous part of a hybrid inclusion), as pointed out in [9], forward invariance of a set is a property that depends on the system's dynamics outside the set. Therefore, in the following results, our flow conditions concern only a neighborhood of the boundary  $\partial K$  relative to the complement of  $K$ .

**Theorem 1** Given a hybrid system  $\mathcal{H} = (C, F, D, G)$  and a  $\mathcal{C}^1$  barrier function candidate  $B$  defining the set  $K$  in (6). Then, the set  $K$  is forward pre-invariant for  $\mathcal{H}$  if, for each  $i \in \{1, 2, \dots, m\}$ , there exists a neighborhood  $U(M_i)$  such that

$$\langle \nabla B_i(x), \eta \rangle \leq 0 \quad \forall x \in (U(M_i) \setminus K_{ei}) \cap C \text{ and} \\ \forall \eta \in F(x) \cap T_C(x), \quad (11)$$

$$B(\eta) \leq 0 \quad \forall \eta \in G(x), \quad \forall x \in D \cap K, \quad (12)$$

$$G(x) \subset C \cup D \quad \forall x \in D \cap K, \quad (13)$$

where  $K_{ei}$  and  $M_i$  are defined in (8)-(9) and  $T_C$  is the contingent cone of the set  $C$ .

**PROOF.** To prove the statement, we proceed by contradiction. Let us assume that (11) and (13) hold and the set  $K$  is not forward pre-invariant for  $\mathcal{H}$ . That is, there exists a maximal solution  $x$  starting from  $x_o \in K$  that leaves the set  $K$  following one of the scenarios (Sc1) and (Sc2). First, suppose that the solution  $x$  leaves the set  $K$  after a jump from  $K$  to  $\mathbb{R}^n \setminus K$  following the scenario (Sc1). This implies, using (13) and the definition of  $B$ , the existence of  $k \in \{1, 2, \dots, m\}$  and  $(t, j) \in \text{dom } x$  such that  $(t, j+1) \in \text{dom } x$  and  $B_k(x(t, j+1)) > 0$  with  $x(t, j+1) \in G(x(t, j))$ . However,  $x(t, j) \in K \cap D$ , hence using (12), it follows that  $B(x(t, j+1)) \leq 0$ ; in fact  $B(\zeta) \leq 0$  for all  $\zeta \in G(x(t, j))$ . The latter fact yields a contradiction. Next, suppose that the solution  $x$  leaves the set  $K$  by flowing under scenario (Sc2). We conclude in this case that there exists  $k \in \{1, 2, \dots, m\}$  such that  $x((t'_1, t'_2), j') \subset (U(\partial K_k) \setminus K_k) \cap C$ , where  $t'_1, t'_2$ , and  $j'$  are as in (Sc2). Next, since the function  $B$  is assumed to be continuously differentiable and the solution  $x(\cdot, j')$  is absolutely continuous on the interval  $[t'_1, t'_2]$ , it follows that  $B(x(\cdot, j'))$  is also absolutely continuous on that interval. By integration, it follows that

$$B_k(x(t'_2, j')) - B_k(x(t'_1, j')) = \\ \int_{t'_1}^{t'_2} \langle \nabla B_k(x(t, j')), \dot{x}(t, j') \rangle dt > 0 \quad (14)$$

since  $B_k(x(t, j')) > 0$  for all  $t \in [t'_1, t'_2]$  and  $B_k(x(t'_1, j')) = 0$ . However,  $x((t'_1, t'_2), j') \subset (U(\partial K) \setminus K) \cap C$  and,

using [30, Lemma 2], we conclude that  $\dot{x}(t, j') \in T_C(x(t, j'))$  for almost all  $t \in [t'_1, t'_2]$ . Moreover, using (11), we conclude that, for almost all  $t \in (t'_1, t'_2)$ ,  $\langle \nabla B_k(x(t, j)), \eta \rangle \leq 0$  for all  $\eta \in F(x(t, j)) \cap T_C(x(t, j))$ . Hence,  $B_k(x(t'_2, j)) - B_k(x(t'_1, j)) \leq 0$ . Hence, the contradiction with (14) follows.  $\square$

**Example 1** Consider the hybrid system  $\mathcal{H}$  with the data

$$\begin{aligned} C &:= \{x \in \mathbb{R}^2 : x_2 \geq 0, x_1 \in [-1, 1]\}, \\ F(x) &:= \begin{bmatrix} -x_2^2 \\ x_2 x_1 - x_2([2, 4] - |x|^2) \end{bmatrix} \quad \forall x \in C, \\ D &:= \{x \in \mathbb{R}^2 : x_2 \leq 0, |x| < 1\}, \\ G(x) &:= [0, x_2] \times [0, |x_1|] \quad \forall x \in D. \end{aligned}$$

We establish forward pre-invariance for the closed set  $K := \{x \in C \cup D : |x|^2 \leq 1, x_2 \geq 0\}$  using Theorem 1. To this end, we note that the set  $K$  can be written as in (6) using the  $C^1$  barrier function candidate  $B(x) = [B_1(x) \ B_2(x)]^\top := [(|x|^2 - 1) \ -x_2]^\top$ . Furthermore,  $D \cap K = (-1, 1) \times \{0\}$  and, for each  $x \in D \cap K$ ,  $G(x) = [0 \ 0, 1]|x_1|]^\top \subset C \cup D$ ; hence, (13) holds. Moreover, for each  $x \in K \cap D$  and for each  $\eta \in G(x)$ , there exists  $\epsilon \in [0, 1]$  such that  $B(\eta) = [(\epsilon|x|^2 - 1) \ -\epsilon|x_1|]^\top \subset \mathbb{R}_{\leq 0} \times \mathbb{R}_{\leq 0}$ ; thus, (12) holds. Next, we note that the set  $(U(M_2) \setminus K_{e2}) \cap C$  is empty and one can choose  $(U(M_1) \setminus K_{e1}) \cap C = \{x \in C : |x| \in (1, 2)\}$ . Consequently, for each  $\eta \in F(x)$ , there exists  $\epsilon \in [0, 2]$  such that  $\langle \nabla B_1(x), \eta \rangle = -x_2^2(2 + \epsilon - |x|^2) \leq 0$  for all  $x \in U(M_1) \setminus K_{e1} \cap C$ . Hence, (11) holds and forward pre-invariance for  $\mathcal{H}$  of the set  $K$  follows. Note that (11) does not hold on the entirety of  $C \setminus K$ .

In the following example, we apply Theorem 1 on a hybrid system including explicit logic variables.

**Example 2 (Thermostat)** Consider the hybrid model of the thermostat system proposed in [33, Example 1.9] with  $x := [q \ z]^\top \in \mathbb{R}^2$ ,

$$\begin{aligned} C &:= (\{0\} \times C_0) \cup (\{1\} \times C_1), \\ C_0 &:= \{z \in \mathbb{R} : z \geq z_{\min}\}, \quad C_1 := \{z \in \mathbb{R} : z \leq z_{\max}\}, \\ F(x) &:= [0 \quad -z + z_o + z_\Delta q]^\top \quad \forall x \in C, \\ D &:= (\{0\} \times D_0) \cup (\{1\} \times D_1), \\ D_0 &:= \{z \in \mathbb{R} : z \leq z_{\min}\}, \quad D_1 := \{z \in \mathbb{R} : z \geq z_{\max}\}, \\ G(x) &:= [1 - q \quad z]^\top \quad \forall x \in D, \end{aligned}$$

where  $z$  is the temperature of the room,  $z_o$  represents the natural temperature of the room when the heater is not used,  $z_\Delta$  the capacity of the heater to raise the temperature in the room by always being on, and  $q$  the state of the heater, which is 1 (on) or 0 (off). We want to keep the temperature between  $z_{\min}$  and  $z_{\max}$  satisfying  $z_o < z_{\min} < z_{\max} < z_o + z_\Delta$ . Using Theorem 1, we will show that the set  $K := \{[q \ z]^\top \in C \cup D : z \in [z_{\min}, z_{\max}]\} =$

$\{0, 1\} \times [z_{\min}, z_{\max}]$  is forward pre-invariant. To do so, we propose the barrier function candidate  $B(x) = [B_1(x) \ B_2(x)]^\top := [z - z_{\max} \ z_{\min} - z]^\top$ . To verify (13), we note that  $C \cup D = \{0, 1\} \times \mathbb{R}$  and  $[1 - q \ z]^\top \in C \cup D$  for all  $[q \ z]^\top \in C \cup D$ . Hence, (13) is satisfied. Moreover,  $B(G(q, z)) = B([(1 - q) \ z]^\top) = B([q \ z]^\top) \leq 0$  for all  $[q \ z]^\top \in K \cap D$ , the latter inequality holds by definition of the barrier candidate  $B$ . Hence, (12) is also satisfied. Finally, to verify (11), we note that  $K_{e1} = \mathbb{R} \times (-\infty, z_{\max}]$ ,  $K_{e2} = \mathbb{R} \times [z_{\min}, +\infty)$ , and  $M_i = \partial K_{ei} \cap (C \cup D)$  for all  $i \in \{1, 2\}$ . Furthermore, for some  $\epsilon > 0$ ,  $(U(M_1) \setminus K_{e1}) \cap C = \{0\} \times (z_{\max}, z_{\max} + \epsilon)$ , and  $(U(M_2) \setminus K_{e2}) \cap C = \{1\} \times (z_{\min} - \epsilon, z_{\min})$ . As a result,  $\langle \nabla B_1(x), F(x) \rangle = z_o - z \leq 0$  for all  $x \in (U(M_1) \setminus K_{e1}) \cap C$ , and  $\langle \nabla B_2(x), F(x) \rangle = z - z_o - z_\Delta \leq 0$  for all  $x \in (U(M_2) \setminus K_{e2}) \cap C$ .

**Remark 4** When the set  $K$  is defined as the zero sublevel set of a scalar barrier function candidate  $B$ ; namely,  $K = \{x \in C \cup D : B(x) \leq 0\}$  with  $m = 1$ , condition (11) in Theorem 1 reduces to

$$\langle \nabla B(x), \eta \rangle \leq 0 \quad \forall x \in (U(\partial K) \setminus K) \cap C \text{ and } \forall \eta \in F(x) \cap T_C(x). \quad (15)$$

**Example 3 (Bouncing ball)** Consider the bouncing ball hybrid model  $\mathcal{H} = (C, F, D, G)$  with  $x \in \mathbb{R}^2$ ,

$$\begin{aligned} F(x) &:= [x_2 \quad -\gamma]^\top \quad \forall x \in C, \\ C &:= \{x \in \mathbb{R}^2 : x_1 > 0, \text{ or } x_1 = 0 \text{ and } x_2 \geq 0\}, \\ G(x) &:= [0 \quad -\lambda x_2]^\top \quad \forall x \in D, \\ D &:= \{x \in \mathbb{R}^2 : x_1 = 0, x_2 \leq 0\}. \end{aligned}$$

The constants  $\gamma > 0$  and  $\lambda \in [0, 1]$  are the gravity acceleration and the restitution coefficient, respectively. Consider the closed set

$$K := \{x \in C \cup D : 2\gamma x_1 + (x_2 - 1)(x_2 + 1) \leq 0\}.$$

The set  $K$  can be seen as the sublevel set where the total energy of the ball is less or equal than 1/2. Hence,  $B(x) := 2\gamma x_1 + (x_2 - 1)(x_2 + 1)$  is a barrier function candidate defining the set  $K$  as in Definition 3. To conclude forward pre-invariance of the set  $K$  using Theorem 1, we start noticing that  $\langle \nabla B(x), F(x) \rangle = 0$  for all  $x \in C$ ; hence, (15) is satisfied. Moreover, for every  $x \in K \cap D$ ,  $B(G(x)) = 2\gamma x_1 + \lambda^2 x_2^2 - 1 \leq 2\gamma x_1 + x_2^2 - 1 \leq 0$  since  $\lambda \in [0, 1]$ . Hence, (12) is satisfied. Finally, (13) is satisfied since  $G(D) = \{0\} \times \mathbb{R}_{\geq 0} \subset C \cup D$ .

**Remark 5** The flow condition (11) in Theorem 1 is more general than those in [17, 18, 20, 23, 24] in the sense that the inequality in (11) does not need to hold on the entire set  $C$ . In [17, 18], the flow condition (11) is expressed as

$$\langle \nabla B(x), \eta \rangle \leq \rho(B(x)) \quad \forall x \in C \quad \forall \eta \in F(x),$$

where  $\rho : \mathbb{R} \rightarrow \mathbb{R}$  is an extended class- $\mathcal{K}$  function; namely,  $\rho(0) = 0$  and  $B \mapsto \rho(B)$  is strictly decreasing. Furthermore, in [23, 24], the function  $\rho : \mathbb{R} \rightarrow \mathbb{R}$  is assumed to be only locally Lipschitz with  $\rho(0) = 0$ . Hence, the function  $\rho$  is allowed to be positive provided that its growth is bounded locally by a linear function. We consider this type of relaxation in Section 3.4. This being said, for continuous-time systems with inputs, when the flow inequalities hold on the entire state space, as shown in [37], numerical methods can be employed to compute an input value that assures safety.

**Remark 6** From conditions (11) in Theorem 1, it is straightforward to conclude that it is enough for each barrier function candidate  $B_i$  to be of class  $C^1$  only on a neighborhood of the boundary  $M_i$ .

**Remark 7** In Theorem 1 (as well as in upcoming results), the jump condition (12) in Theorem 1 can be formulated using a different barrier function candidate than the one used to formulate the flow condition (11) in Theorem 1. However, the two different barrier function candidates still need to define the same set  $K$ , according to (6). In this paper, for simplicity, we present results using the same barrier function candidate in the flow and the jump conditions, but extensions to the case where they are different are straightforward.

### 3.2 Pre-Invariance Under Lipschitz-Like Flow Map

In Theorem 1, the inequality in (11) needs to be satisfied on a neighborhood outside the set  $K$  rather than just on  $\partial K$ . This is also the case for  $m = 1$  for which (15) is required; see Remark 4. To assess the possibility of relaxing such requirement, we consider the flow condition

$$\langle \nabla B_i(x), \eta \rangle \leq 0 \quad \forall x \in M_i \cap C, \quad \forall \eta \in F(x). \quad (16)$$

We will show that under some regularity assumptions on  $F$  and  $B$ , condition (16) can be used to conclude forward pre-invariance of the set  $K \subset C \cup D$ . Furthermore, as we show in Example 5, when we further relax (16) to

$$\langle \nabla B_i(x), \eta \rangle \leq 0 \quad \forall x \in M_i \cap C, \quad \forall \eta \in F(x) \cap T_C(x),$$

we fail to conclude forward pre-invariance of the set  $K$ .

When (16) and (12)-(13) hold, the closed set  $K \subset C \cup D$  can fail to be forward pre-invariant for the reasons enumerated below, where we assume without loss of generality that  $m = 1$  (i.e.,  $B$  is a scalar function).

- 1) Assume the existence of  $x_o \in \partial K \cap \text{int}(C)$  such that  $\nabla B(x_o) = 0$ . Assume also that  $F(x_o) \subset D_{\mathbb{R}^n \setminus K}(x_o)$ , where  $D_{\mathbb{R}^n \setminus K}$  is the Dubovitsky-Miliutin cone of the set  $\mathbb{R}^n \setminus K$ . In this case, condition (16) is satisfied at

$x_o$ . However, according to [30, Theorem 6], there exists a nontrivial solution starting from  $x_o$  and flowing outside the set  $K$ . Hence, the set  $K$  is not forward pre-invariant although (16) is satisfied; cf. [17, Proposition 1]. To handle this situation, one needs to assume that the gradient of  $B$  is non-degenerate on  $\partial K_e \cap C$ ; namely,

$$\nabla B(x) \neq 0 \quad \forall x \in \partial K_e \cap C. \quad (17)$$

- 2) When the solutions starting from  $x_o \in \partial K \cap C$  are nonunique, even if  $\nabla B(x_o) \neq 0$ , when  $\langle \nabla B(x_o), \eta \rangle = 0$  for any  $\eta \in F(x_o)$ , we can always consider the existence of a solution starting from  $x_o \in \partial K \cap C$  with a speed that is tangent to  $\partial K$  but leaving the set  $K$ . Such a scenario is illustrated in Example 4 below. We also notice that this pathology does not occur when  $\dot{x} \in F(x)$  for all  $x \in C$  has unique solutions.

The following example is inspired by [8, Page 1751].

**Example 4** Consider the two-dimensional differential equation  $\dot{x} = [1 \quad \sqrt{|x_2|}]^\top =: F(x)$  and the set  $K := \{x \in \mathbb{R}^2 : x_2 \leq 0\}$ . Note that this system can be interpreted as a hybrid system with  $C = \mathbb{R}^2$ ,  $D$  empty, and  $G$  arbitrary. The set  $K$  can be defined using the barrier function candidate  $B(x) := x_2$  satisfying  $|\nabla B(x)| = 1 \neq 0$  and  $\langle \nabla B(x), F(x) \rangle = 0$  for all  $x \in \partial K$ . However, the set  $K$  is not forward pre-invariant since  $x(t) = [t \quad (1/4)t^2]^\top$  defined for all  $t \geq 0$  is a solution starting from  $x(0) = 0 \in K$  that leaves the set  $K$ .

The latter example confirms the fact that, for differential inclusions, forward pre-invariance of a closed set  $K$  is a property of the system outside the set  $K$  rather than on its boundary or in its interior. However, when the flow map  $F$  satisfies extra regularity conditions outside the set  $K$ , it is possible to restrict the conditions in (11) and (15) to hold only on the boundary  $\partial K$ . This is possible, for example when  $F$  is locally Lipschitz as shown in [13, 38, 39] for differential equations and in [9] for differential inclusions. In the aforementioned references, contingent-cone-based conditions are used and shown to be necessary as well as sufficient, provided that the system's dynamics is defined on an open set containing the closed set  $K$ . For differential equations defined in  $\mathbb{R}^n$  with locally Lipschitz right-hand side, the latter contingent-cone-based conditions are expressed in terms of a scalar barrier function candidate in [17].

On the other hand, to conclude forward pre-invariance using flow conditions satisfied only on the boundary of the set  $K$ , the Lipschitz regularity of the flow map  $F$  can be relaxed by modifying the right-hand side in [30, Definition 15] using uniqueness functions. The latter is shown in [40] for differential equations.

**Definition 4 (Uniqueness function)** A function  $\rho : \mathbb{R} \rightarrow \mathbb{R}$  is said to be a uniqueness function if, for each

continuous function  $\delta : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$  such that  $\delta(0) = 0$  and there exists  $\epsilon > 0$  such that

$$\limsup_{h \rightarrow 0^+} \frac{\delta(t+h) - \delta(t)}{h} \leq \rho(\delta(t)) \text{ for a.a. } t \in [0, \epsilon], \quad (18)$$

it follows that  $\delta(t) = 0$  for all  $t \in [0, \epsilon]$ .

**Remark 8** Uniqueness functions, in this paper, are introduced in a slightly different way compared to [40]. Indeed, in this reference, a function  $\rho : \mathbb{R} \rightarrow \mathbb{R}$  is said to be a uniqueness function if, for each continuous function  $\delta : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$  such that  $\delta(0) = 0$  and there exists  $\epsilon > 0$  such that, in addition to (18), we have

$$\limsup_{h \rightarrow 0^+} \frac{\delta(t) - \delta(t-h)}{h} \leq \rho(\delta(t)) \text{ for a.a. } t \in [0, \epsilon],$$

it follows that  $\delta(t) = 0$  for all  $t \in [0, \epsilon]$ . That is, the notion of uniqueness function used in this paper considers only the upper-right Dini derivative.

In the following statement, we propose flow and jump conditions that are sufficient and need to be satisfied only on elements of the set  $K$  provided that the following assumption holds.

**Assumption 1** For every  $x \in \partial K_e \cap C$ ,

$$\begin{aligned} \exists v \in \mathbb{R}^n : \langle \nabla B_i(x), v \rangle < 0 \\ \forall i \in \{1, 2, \dots, m\} \text{ s.t. } B_i(x) = 0. \end{aligned} \quad (19)$$

Assumption 1 is known as *transversality condition* in [32] and allows to define the contingent cone  $T_K$  at the intersection between different zero-level sets — see [30, Lemma 3]. Furthermore, Assumption 1 reduces to (17) when the barrier function  $B$  is scalar.

**Theorem 2** Given a hybrid system  $\mathcal{H} = (C, F, D, G)$  and a  $C^1$  barrier function candidate  $B$  defining the set  $K$  in (6). Furthermore, assume that there exists a neighborhood  $U(\partial K)$  such that, for each  $x \in (U(\partial K) \setminus K) \cap C$  and for each  $y \in \partial(K \cap C)$ ,

$$(x-y)^\top F(x) \subset (x-y)^\top F(y) + |x-y|\rho(|x-y|)\mathbb{B} \quad (20)$$

with  $\rho : \mathbb{R} \rightarrow \mathbb{R}$  a uniqueness function. Then, the closed set  $K$  is forward pre-invariant provided that (12)-(13), (16), and Assumption 1 hold, and there exists a neighborhood  $U(\partial K_e \cap \partial C)$  such that one of the following extra conditions holds:

(a) For any  $x \in (U(\partial K_e \cap \partial C) \cap \partial K_e) \setminus C$ , (19) holds and

$$\begin{aligned} \langle \nabla B_i(x), \eta \rangle \leq 0 \quad \forall \eta \in F(x) \text{ and} \\ \forall i \in \{1, 2, \dots, m\} \text{ s.t. } B_i(x) = 0. \end{aligned} \quad (21)$$

(b) For any  $x \in U(\partial K_e \cap \partial C) \cap \partial K \cap \partial C$ ,

$$F(x) \subset T_{K \cap C}(x). \quad (22)$$

(c) The set  $C$  is convex and (22) holds for all  $x \in \partial K_e \cap \partial C$ .

**PROOF.** The proof that the solutions starting from the set  $K$  cannot jump outside according to (Sc1), under (12)-(13), is the same as in the proof of Theorem 1. Next, we prove that the solutions starting from  $\partial K_e \cap C$  cannot leave the set  $K$  by flowing as in scenario (Sc2). For this purpose, we adapt the steps of the proof presented in [40, Proof of Theorem 1] to our more general setting. Let  $t'_1 \geq 0$  and  $t'_2 > t'_1$  be such that there exists a solution flowing from  $x_o := x(t'_1, 0) \in \partial K \cap C$  and satisfying  $x(t, 0) \in (U(\partial K) \setminus K) \cap C$  for all  $t \in (t'_1, t'_2)$ . We use  $y_1$  to denote the projection of  $x(t, 0)$  on the set  $K_e$  and  $y_2$  to denote the projection of  $x(t, 0)$  on the set  $K \cap C$ . Furthermore, we define  $\delta_1(t) := |x(t, 0) - y_1|$  and  $\delta_2(t) := |x(t, 0) - y_2|$ . It follows that  $\delta_i(t'_1) = 0$  and  $\delta_i(t) > 0$  for all  $t \in (t'_1, t'_2)$ , since  $x(t, 0) \in (U(\partial K) \setminus K) \cap C$  for all  $t \in (t'_1, t'_2)$ , for each  $i \in \{1, 2\}$ . Using the identity  $a - b = (a^2 - b^2)/(a + b)$  for  $a$  and  $b$  nonnegative and for any  $h > 0$  such that  $t$  and  $t + h$  in  $(t'_1, t'_2)$ , we derive the inequality

$$\begin{aligned} \delta_i(t+h) - \delta_i(t) &\leq |x(t+h, 0) - y_i| - |x(t, 0) - y_i| \\ &= \frac{|x(t+h, 0) - y_i|^2 - |x(t, 0) - y_i|^2}{|x(t+h, 0) - y_i| + |x(t, 0) - y_i|} \end{aligned} \quad (23)$$

for each  $i \in \{1, 2\}$ . Furthermore, for almost all  $t \in (t'_1, t'_2)$ , we replace  $x(t+h, 0)$  by

$$x(t+h, 0) = x(t, 0) + h\dot{x}(t, 0) + o(h) \quad (24)$$

with  $o(h)$  the remainder of the first order Taylor expansion of  $h \mapsto x(t+h)$  around  $h = 0$ , which satisfies  $\lim_{h \rightarrow 0} o(h)/h = 0$ . Using the previous limit and the inequality

$$|x(t+h, 0) - y_i| \leq |x(t, 0) - y_i| + h|\dot{x}(t, 0)| + o(h)/h,$$

we obtain that, for all  $i \in \{1, 2\}$  and for almost all  $t \in (t'_1, t'_2)$ ,

$$\limsup_{h \rightarrow 0^+} \frac{\delta_i(t+h) - \delta_i(t)}{h} \leq \frac{(x(t, 0) - y_i)^\top \dot{x}(t, 0)}{|x(t, 0) - y_i|}. \quad (25)$$

Next, we have the following claim:

**Claim 1** Under (16) and (19), the following is true.

(cl1) If (a) holds, then  $(x(t, 0) - y_1)^\top \eta_y \leq 0$  for all  $\eta_y \in F(y_1)$ .



(cl2) If (b) or (c) hold, then  $(x(t, 0) - y_2)^\top \eta_y \leq 0$  for all  $\eta_y \in F(y_2)$ .

To prove the claim, we proceed as follows:

- Under (a) in Theorem 2, we show that  $F(y_1) \subset T_{K_e}(y_1)$ . Indeed, under (16) and (19) and using [30, Lemma 3], it follows that  $F(y_1) \subset T_{K_e}(y_1)$  when  $y_1 \in \partial K_e \cap C$ . Similarly, using the same argument under (a), when  $y_1 \in (U(\partial K_e \cap \partial C) \cap \partial K_e) \setminus C$ , we also have  $F(y_1) \subset T_{K_e}(y_1)$ . Finally, we use the fact that when  $t'_2$  is sufficiently small, the element  $y_1$  corresponding to the projection of  $x(t, 0) \in (U(\partial K) \setminus K) \cap C$  on the set  $K_e$  belongs necessarily to either  $\partial K_e \cap C$  or  $(U(\partial K_e \cap \partial C) \cap \partial K_e) \setminus C$ .
- Under (b) or (c) in Theorem 2, we show that  $F(y_2) \subset T_{K \cap C}(y_2)$ . Indeed, under (16), Assumption 1, and using [30, Lemma 3], it follows that  $F(y_2) \subset T_{K_e}(y_2) = T_{K \cap C}(y_2)$  when  $y_2 \in \partial K_e \cap \text{int}(C)$ . Next, under (b), when  $y_2 \in U(\partial K_e \cap \partial C) \cap \partial K \cap \partial C$ , we also have  $F(y_2) \subset T_{K \cap C}(y_2)$ . Finally, we use the fact that when  $x_o \in \partial K_e \cap C$  and for  $t'_2 > 0$  sufficiently small, the element  $y_2$  corresponding to the projection of  $x(t, 0)$  on the set  $K \cap C$  belongs necessarily to either  $(\partial K_e \cap \text{int}(C))$  or  $(U(\partial K_e \cap \partial C) \cap \partial K \cap \partial C)$ . Furthermore, under (c), the set  $C$  is convex and we show that  $y_2 \in \partial K_e \cap C$ . Hence, it becomes enough to have  $F(y_2) \subset T_{K \cap C}(y_2)$  when  $y_2 \in \partial K_e \cap \partial C$ , which is true under (c). Finally, to show that  $y_2 \in \partial K_e \cap C$  when  $C$  is convex, we use the fact that  $y_2$  corresponds to a projection on the set  $K \cap C$ . Hence, either  $y_2 \in \partial K_e \cap C$  or  $y_2 \in \partial C \cap \text{int}(K_e)$ . We propose to exclude the latter case using contradiction. That is, assume that  $y_2 \in \partial C \cap \text{int}(K_e)$  and consider the line segment relating  $y_2$  to  $x(t, 0)$  denoted by  $[y_2, x(t, 0)]$ . Since both  $y_2$  and  $x(t, 0)$  lie in the set  $C$  and since the set  $C$  is convex it follows that the segment  $[y_2, x(t, 0)]$  also belongs to  $C$ . Furthermore, since  $x(t, 0) \in C \setminus K_e$  and  $y_2 \in \text{int}(K_e)$  it follows the existence of  $y_o$  belonging to the open segment  $(x(t, 0), y_2)$  such that  $y_o \in \partial K_e \cap C = \partial K \cap C$ . Hence  $|x(t, 0) - y_o| < |x(t, 0) - y_2| = \min_{z \in K \cap C} \{|x(t, 0) - z|\}$ , which yields to a contradiction.

Now, to conclude (cl1) and (cl2), we use the inequalities

$$\begin{aligned} |x(t, 0) - y_1| &\leq |x(t, 0) - y_1 - h\eta_y| + |y_1 + h\eta_{y_1}|_{K_e}, \\ |x(t, 0) - y_2| &\leq |x(t, 0) - y_2 - h\eta_y| + |y_2 + h\eta_{y_2}|_{K \cap C}, \end{aligned}$$

where  $(\eta_{y_1}, \eta_{y_2}) \in F(y_1) \times F(y_2)$ . To obtain the previous inequalities, we used the fact that the functions  $|\cdot|_{K_e}$  and  $|\cdot|_{K \cap C}$  are globally Lipschitz with a Lipschitz constant equal to 1,  $|x(t, 0) - y_1| = |x(t, 0)|_{K_e}$ , and  $|x(t, 0) - y_2| = |x(t, 0)|_{K \cap C}$ . Next, by taking the square in both sides of the last two inequalities, and dividing by  $h$ , we obtain,

for each  $\eta_{y_1} \in F(y_1)$ ,

$$\begin{aligned} |x(t, 0) - y_1|^2/h &\leq |x(t, 0) - y_1 - h\eta_{y_1}|^2/h + \\ &|y_1 + h\eta_{y_1}|_{K_e}^2/h + \\ &2|x(t, 0) - y_1 - h\eta_{y_1}||y_1 + h\eta_{y_1}|_{K_e}/h, \end{aligned}$$

which implies that

$$\begin{aligned} |x(t, 0) - y_1|^2/h &\leq |x(t, 0) - y_1|^2/h + h|\eta_{y_1}|^2 - \\ &2(x(t, 0) - y_1)\eta_{y_1} + h(|y_1 + h\eta_{y_1}|_{K_e}/h)^2 + \\ &2|x(t, 0) - y_1 - h\eta_{y_1}||y_1 + h\eta_{y_1}|_{K_e}/h. \end{aligned}$$

Similarly, for each  $\eta_{y_2} \in F(y_2)$ ,

$$\begin{aligned} |x(t, 0) - y_2|^2/h &\leq |x(t, 0) - y_2|^2/h + h|\eta_{y_2}|^2 - \\ &2(x(t, 0) - y_2)\eta_{y_2} + \\ &h(|y_2 + h\eta_{y_2}|_{K \cap C}/h)^2 + \\ &2|x(t, 0) - y_2 - h\eta_{y_2}||y_2 + h\eta_{y_2}|_{K \cap C}/h. \end{aligned}$$

Finally, letting  $h \rightarrow 0^+$  through a suitable sequence, (cl1) and (cl2) are proved using the fact that  $\liminf_{h \rightarrow 0^+} |y_1 + h\eta_{y_1}|_{K_e}/h = 0$  and  $\liminf_{h \rightarrow 0^+} |y_2 + h\eta_{y_2}|_{K \cap C}/h = 0$  since we already showed that  $\eta_{y_1} \in T_{K_e}(y_1)$  under (a) in Theorem 2 and  $\eta_{y_2} \in T_{K \cap C}(y_2)$  under (b) or (c) in Theorem 2.

Using the Claim 1, for each  $\eta_y \in F(y_i)$ , the term  $-\frac{(x(t, 0) - y_i)^\top \eta_y}{|x(t, 0) - y_i|}$  can be added in (25) which then can be rewritten as

$$\limsup_{h \rightarrow 0^+} \frac{\delta_i(t+h) - \delta_i(t)}{h} \leq \frac{(x(t, 0) - y_i)^\top (\dot{x}(t, 0) - \eta_y)}{|x(t, 0) - y_i|}, \quad (26)$$

where  $i = 1$  if (a) in Theorem 2 holds and  $i = 2$  if (b) or (c) in Theorem 2 hold. Since  $\dot{x}(t, 0) \in F(x(t, 0))$  for almost all  $t \in (t'_1, t'_2)$ , using (20) with  $x = x(t, 0)$  and  $y = y_i$ , it is always possible to find  $\eta_y^* \in F(y_i)$  such that

$$(x(t, 0) - y_i)^\top (\dot{x}(t, 0) - \eta_y^*)^\top \leq |x(t, 0) - y_i| \rho(|x(t, 0) - y_i|).$$

Applying this inequality to (26) and replacing  $\eta_y$  therein by  $\eta_y^*$ , we obtain

$$\limsup_{h \rightarrow 0^+} \frac{\delta_i(t+h) - \delta_i(t)}{h} \leq \rho(|x(t, 0) - y_i|) = \rho(\delta_i(t))$$

for almost all  $t \in (t'_1, t'_2)$  with  $i = 1$  if (a) holds and  $i = 2$  (b) or (c) hold. Since the function  $\rho$  is a uniqueness function, it follows that  $\delta_i(t) = 0$  for all  $t \in (t'_1, t'_2)$ . Hence, the contradiction follows.  $\square$




Fig. 1. Illustration of the system in Example 5.

**Remark 9** Note that condition (20) holds for free when  $F$  is locally Lipschitz. In fact, every locally Lipschitz map  $F$  satisfies (20) with  $\rho(\omega) = k\omega$  for some  $k > 0$ . Condition (20) is more general than Lipschitzness and allows for functions  $\rho$  that are not necessarily linear. In particular, when  $F$  is such that (20) holds for  $\rho(\omega) := \omega \log \omega$ , then  $\rho$  is a uniqueness function [41]. The latter function belongs to the more general class of Osgood functions that are uniqueness functions but not necessarily locally Lipschitz [40].

**Remark 10** The flow condition (21) in Theorem 2 is a reinterpretation, in terms of barrier functions, of the well-known contingent cone-based condition used in [9, 13, 38, 39] provided that Assumption 1 holds.

In Example 5, we show that when none of the conditions (a)-(c) in Theorem 2 is satisfied, there exist situations where  $K$  fails to be forward pre-invariant. Also, we show that the aforementioned conditions are only sufficient.

**Example 5** Consider the differential inclusion  $\mathcal{H} = (C, F, \emptyset, \star)$ , where  $F(x) := [1 \ 0]^\top$  for all  $x \in C$  and

$$C := \{x \in \mathbb{R}^2 : |x_2| \geq x_1^2\} \cup \{x \in \mathbb{R}^2 : x_1 \leq 0\} \cup \{x \in \mathbb{R}^2 : x_2 = 0\}.$$

Furthermore, consider the barrier function candidate

$$B(x) := \begin{cases} x_2 & \text{if } x_1 \leq 0 \\ x_2 + x_1^3 & \text{otherwise} \end{cases}$$

defining a closed set  $K$  according to (6). We will show that (16)-(17) are satisfied; however, none of the conditions (a)-(c) in Theorem 2 is satisfied. As a result, we show that the set  $K$  is not forward pre-invariant. Indeed, note that  $\nabla B(x) = [0 \ 1]^\top$  if  $x_1 \leq 0$  and  $\nabla B(x) = [2x_1^2 \ 1]^\top$  otherwise; hence, (17) is satisfied. Furthermore, for each  $x \in \partial K_e \cap C = \{x \in \mathbb{R}^2 : x_2 = 0, x_1 \leq 0\}$ , we have  $\langle \nabla B(x), F(x) \rangle = \langle [0 \ 1]^\top, [1 \ 0]^\top \rangle \leq 0$ ; hence, (16) is satisfied. Moreover, (12)-(13) are trivially satisfied since  $D = \emptyset$ . Next, it is easy to see that the set  $C$  is not convex; hence, condition (c) is not satisfied. Furthermore, when  $x \in (U(\partial K_e \cap \partial C) \cap \partial K_e) \setminus C = \{x \in \mathbb{R}^2 : x_2 = -x_1^3, x_1 \in (0, \epsilon], \epsilon > 0\}$ ,  $\langle \nabla B(x), F(x) \rangle =$

$\langle [2x_1^2 \ 1]^\top, [1 \ 0]^\top \rangle = 2x_1^2 > 0$ ; hence, (a) is not satisfied. Next, for any  $x \in U(\partial K_e \cap \partial C) \cap \partial K \cap \partial C = \{x \in \mathbb{R}^2 : x_2 = -x_1^3, x_1 \in [0, \epsilon], \epsilon > 0\}$ , if  $x = 0$ , then  $F(0) \in T_{K \cap C}(0)$ ; however, if  $x \neq 0$ , then  $F(x) \notin T_C(x)$  because  $F(x)$  points outside the set  $\{x \in \mathbb{R}^2 : |x_2| \geq x_1^2\}$ , which defines  $C$  when  $x_1 > 0$ . Hence, (b) is also not satisfied. Finally, the constrained differential inclusion  $\mathcal{H} = (C, F, \emptyset, \star)$  admits the solution  $x(t) = [t \ 0]^\top, t \geq 0$ , starting from  $x_0 = 0 \in K$ , that leaves the set  $K$ .

Now, in order to show that none of the conditions (a)-(c) is necessary, we slightly modify the set  $C$  in order to render the set  $K$  forward pre-invariant while maintaining (a)-(c) unsatisfied. That is, consider the new flow set

$$C_1 := \{x \in \mathbb{R}^2 : |x_2| \geq x_1^2\} \cup \{x \in \mathbb{R}^2 : x_1 \leq 0\}.$$

Note that the unique solution starting from each  $x_0 := [x_{o1} \ x_{o2}]^\top \in K$  satisfies  $x(t) = [x_{o1} + t \ x_{o2}]^\top$  for all  $t \geq 0$  and cannot leave the set  $K$  while remaining in  $C$ . However, using the same arguments as in the previous paragraph, we conclude that conditions (a)-(c) remain unsatisfied.

**Example 6** Consider the hybrid system  $\mathcal{H}$  with the data

$$C := \{x \in \mathbb{R}^2 : |x_1| \leq 1\}, \quad F(x) := \begin{bmatrix} 1 \\ -[0, x_2 \log |x_2|] \end{bmatrix},$$

$$D := \{x \in \mathbb{R}^2 : x_1 = 1\}, \quad G(x) := [-1 \ x_2]^\top.$$

To conclude forward pre-invariance of the set  $K := \{x \in C \cup D : x_2 \leq 0\}$  admitting the  $C^1$  barrier function candidate  $B(x) := x_2$ , we start noting that conditions (12)-(13) are satisfied since  $G(x) \in C$  for all  $x \in D$  and  $B(G(x)) = B(x) \leq 0$  for all  $x \in K$ . Furthermore, conditions (16)-(17) are also satisfied since  $\nabla B(x) = [0 \ 1]^\top$  for all  $x \in \mathbb{R}^2$  and  $\langle \nabla B(x), F(x) \rangle = -[0, x_2 \log |x_2|] \leq 0$  for all  $x \in \partial K_e$ . For the same reason, condition (a) in Theorem 2 is also satisfied. Finally, it remains to show that for any  $x \in (U(\partial K) \setminus K) \cap C$  and for any  $y \in \partial(K \cap C)$ , (20) is satisfied. Indeed, for  $y := [y_1 \ y_2]^\top \in \partial K \cap C$ , it follows that  $y_2 = 0$ . At the same time, when  $x \in (U(\partial K) \setminus K) \cap C$ , it follows that  $x_2 > 0$  which can be chosen sufficiently small so that

$$\begin{aligned} (x - y)^\top F(x) &= (x_1 - y_1) + [0, 1]x_2^2 \log(|x_2|) \\ &\subset (x_1 - y_1) + [-1, 1]x_2^2 \log(|x_2|) \\ &= (x_1 - y_1) + [-1, 1](x_2 - y_2)^2 \log(|x_2 - y_2|) \\ &\subset (x - y)^\top F(y) + [-1, 1]|x - y|^2 \log(|x - y|) \\ &= (x - y)^\top F(y) + [-1, 1]|x - y|\rho(|x - y|), \end{aligned}$$

where  $\rho(w) := w \log w$ , for all  $w \geq 0$ . The function  $\rho$  is Osgood [40]; hence, a uniqueness function.

### 3.3 Pre-Invariance Using Locally Lipschitz Barrier Functions

Another approach to conclude forward pre-invariance using cone conditions without restricting the regularity

of the flow map, consists of replacing the flow condition (11) in Theorem 1 by a cone condition to be satisfied on the external part of a sufficiently small neighborhood of the set  $\partial K_e \cap C$ . Inspired by [9, Theorem 5.2.1], the flow condition in the following statement uses the external contingent cone  $E_K$  defined in (4).

**Theorem 3** *Given a hybrid system  $\mathcal{H} = (C, F, D, G)$  and a barrier function candidate  $B$  defining the set  $K$  in (6). The set  $K$  is forward pre-invariant if (12)-(13) hold, and there exists a neighborhood  $U(\partial K)$  such that*

$$\eta \in E_K(x) \quad \forall \eta \in F(x) \cap T_C(x) \text{ and} \quad (27)$$

$$\forall x \in (U(\partial K) \setminus K) \cap C,$$

where  $E_K$  is introduced in (4).

**PROOF.** The proof that the solutions, under (12)-(13), cannot leave the set  $K$  after a jump according to scenario (Sc1) is the same as in the proof of Proposition 1. Next, as in [9, Theorem 5.2.1], we show that the trajectories starting from the set  $\partial K \cap C$  cannot leave the set  $K$  by flowing according to scenario (Sc2). Indeed, assume that a solution  $x$  starting from  $\partial K \cap C$  leaves the set  $K$  by flowing according to scenario (Sc2). Then, there exists  $j \in \mathbb{N}$ , such that, for  $t'_2$  small enough and for all  $t \in (t'_1, t'_2]$ ,  $x(t, j) \in (U(\partial K) \setminus K) \cap C$ , where  $t'_1$  and  $t'_2$  are as in (Sc2). Furthermore, using [30, Lemma 2], we conclude that  $\dot{x}(t, j) \in T_C(x(t, j))$  for almost all  $t \in [t'_1, t'_2]$ . Next, since the distance function with respect to the set  $K$  is locally Lipschitz and the solution  $x(\cdot, j)$  is absolutely continuous on the interval  $[t'_1, t'_2]$ , it follows that  $\delta_K(\cdot) := |(x(\cdot, j))|_K$  is also absolutely continuous on that same interval. Hence, for almost all  $t \in [t'_1, t'_2]$ , the time derivative  $\dot{\delta}_K(t)$  exists and satisfies

$$\lim_{h \rightarrow 0^+} \frac{\delta_K(t+h) - \delta_K(t)}{h} =$$

$$\liminf_{h \rightarrow 0^+} \frac{|x(t, j) + h\dot{x}(t, j)|_K - |x(t, j)|_K}{h}.$$

Since  $\dot{x}(t, j) \in F(x(t, j)) \cap T_C(x(t, j))$  for almost all  $t \in [t'_1, t'_2]$  and  $x((t'_1, t'_2], j) \subset (U(\partial K) \setminus K) \cap C$ , using (27), we conclude that  $\dot{x}(t, j) \in E_K(x(t, j))$  for almost all  $t \in (t'_1, t'_2)$ , which implies that  $\dot{\delta}_K(t) \leq 0$  for almost all  $t \in (t'_1, t'_2)$ . Thus,  $|x(t'_2, j)|_K - |x(t'_1, j)|_K \leq 0$ . The contradiction follows since  $x(t'_2, j) \in C \setminus K$  and  $x(t'_1, j) \in K$  which means that  $|x(t'_2, j)|_K - |x(t'_1, j)|_K > 0$ .  $\square$

**Remark 11** *The flow condition (27) in Theorem 3 is similar to the flow condition (15) in Remark 4. Indeed, in (27), we are using the distance function  $B(x) := |x|_K$  as a barrier function candidate defining the set  $K$  according to Definition 3. However, since the distance function to a set is only locally Lipschitz, the gradient-based inequality in (15) is replaced by the limit in (4) that has the same*

*implication on the monotonic behavior of  $t \mapsto B(x(t, 0))$  on the neighborhood  $(U(\partial K) \setminus K) \cap C$ . Furthermore, according to [9, Corollary 5.2.3], the external cone condition (27) in Theorem 3 is satisfied provided that, for any  $x \in (U(\partial K) \setminus K) \cap C$ ,  $F(x) \cap T_C(x) \subset F(y) \subset T_K(y)$ , where  $y$  is the projection of  $x$  on the set  $K$ .*

Inspired by the discussion in Remark 11, in the following statement, we replace the flow condition (27) in Theorem 3 by a condition involving a general locally Lipschitz barrier function candidate instead of only the distance function. The proof is in [30].

**Theorem 4** *Given a hybrid system  $\mathcal{H} = (C, F, D, G)$  and a scalar locally Lipschitz barrier function candidate  $B$  defining the set  $K$  in (6). The set  $K$  is forward pre-invariant if (12)-(13) hold, and there exists a neighborhood  $U(\partial K)$  such that*

$$\max_{\zeta \in \partial_C B(x)} \langle \zeta, \eta \rangle \leq 0 \quad \forall \eta \in F(x) \cap T_C(x) \text{ and} \quad (28)$$

$$\forall x \in (U(\partial K) \setminus K) \cap C,$$

where  $\partial_C B$  is the Clarke generalized gradient of  $B$ , see [30, Definition 16].

**Remark 12** *In Theorem 4, we consider only the case of scalar barrier function candidates defining the set  $K$ . Conveniently, when the set  $K$  is defined via multiple locally Lipschitz candidates according to (6), we can then use (10) to construct a scalar barrier function that is locally Lipschitz and at the same time defines the set  $K$  according to (6). Furthermore, if we compare conditions (11) and (28) while replacing the scalar function  $B$  in (28) by the function  $\bar{B}$  in (10), we notice that the inequality in (11), for each  $i \in \{1, 2, \dots, m\}$ , is checked in more points compared to (28). Finally, when the set  $C$  is open and  $K \subset C$ , condition (30) covers what is proposed in [18, Theorem 2] for unconstrained differential inclusions.*

**Example 7** *Consider the hybrid system  $\mathcal{H} = (C, F, D, G)$  with  $C$  and  $G$  as in Example 1, and  $F$  and  $D$  given by*

$$F(x) := \begin{bmatrix} -x_2^2 x_1 \\ -(|x|^2 - [0, 1])(2 - |x|^2) \end{bmatrix} \quad \forall x \in C,$$

$$D := \{x \in \mathbb{R}^2 : x_2 = 0, |x| \leq 1\}.$$

*We employ Theorem 3 to verify forward pre-invariance of the closed set  $K := \{x \in C \cup D : |x|^2 - 1 \leq 0\}$ . Indeed, this set admits the scalar barrier function candidate  $B(x) := x_2(|x|^2 - 1)$ . Furthermore, for any  $x \in D \cap K$ ,  $G(x) \subset C \cup D$ ; hence, (13) is satisfied. Moreover, for every  $x \in C \cap D$  and for every  $\eta \in G(x)$ , there exists  $\alpha \in [0, 1]$  such that  $\eta = [0 \ \alpha|x_1|]^\top$  and  $B(\eta) = \alpha|x_1|(\alpha^2|x_1|^2 - 1)$ ; hence, since  $x_1 \leq 1$  for all  $x \in K \cap D$ , (12) follows. Furthermore, the set  $(U(\partial K) \setminus K) \cap C$  can be chosen to be the open set*

$(U(\partial K) \setminus K) \cap C = \{x \in C : |x_1| < 1, |x| \in (1, 2)\}$ . Note that, the function  $|\cdot|_K$  satisfies  $|x|_K = |x| - 1$  for all  $x \in (U(\partial K) \setminus K) \cap C$  and is differentiable on the latter set. Hence,  $\liminf_{h \rightarrow 0^+} \frac{|x+hv|_K - |x|_K}{h} = \langle [x_1 \ x_2]^\top, v \rangle / |x|$  and  $E_K(x) = \{v \in \mathbb{R}^2 : \langle [x_1 \ x_2]^\top, v \rangle \leq 0\}$  for all  $x \in (U(\partial K) \setminus K) \cap C$ . This implies that (27) holds since, for every  $\eta \in F(x)$  and for every  $x \in (U(\partial K) \setminus K) \cap C$ ,  $\langle [x_1 \ x_2]^\top, \eta \rangle = -x_1^2 x_2^2 - x_2(|x|^2 - \epsilon)(2 - |x|^2) \leq 0$  for some  $\epsilon \in [0, 1]$ .

### 3.4 Pre-Invariance Using a Relaxed Flow Condition

Inspired by the property of the uniqueness functions in Definition 4, we are able to relax the sign of the inequality in (11) provided that a growth condition involving a uniqueness function and the barrier function candidate is satisfied. Such a relaxation follows the lines of what is proposed in [23, 24] for hybrid automata, see Remark 5.

**Proposition 1** *Given a hybrid system  $\mathcal{H} = (C, F, D, G)$  and a  $\mathcal{C}^1$  barrier function candidate  $B$  defining the set  $K$  in (6). The set  $K$  is forward pre-invariant if (12)-(13) hold and there exists a neighborhood  $U(M_i)$  such that*

$$\langle \nabla B_i(x), \eta \rangle \leq \rho(B_i(x)) \quad \forall x \in (U(M_i) \setminus K_{ei}) \cap C \text{ and} \\ \forall \eta \in F(x) \cap T_C(x), \quad (29)$$

where  $\rho : \mathbb{R} \rightarrow \mathbb{R}$  is a uniqueness function.

**PROOF.** Under (12)-(13), the proof that the solutions starting from the set  $K$  cannot jump outside the set  $K$  following scenario (Sc1) is the same as in the proof of Theorem 1. The only remaining way to leave the set  $K$  is by flowing according to scenario (Sc2). We conclude in this case, for  $t'_2$  small enough, the existence of  $k \in \{1, \dots, m\}$  such that  $B_k(x(t, j)) > 0$  for all  $t \in (t'_1, t'_2]$  and  $x((t'_1, t'_2], j) \subset (U(\partial K_k) \setminus K_k) \cap C$ , where  $t'_1$  and  $t'_2$  are as in (Sc2). Furthermore, using [30, Lemma 2], we conclude that  $\dot{x}(t, j) \in T_C(x(t, j))$  for almost all  $t \in [t'_1, t'_2]$ . Hence, using (29), we conclude that

$$\frac{dB_k(x(t, j))}{dt} \leq \rho(B_k(x(t, j))) \quad \text{for a. a } t \in [t'_1, t'_2]$$

with  $B_k(x(t'_1, j)) = 0$ . Since  $\rho$  is a uniqueness function, we conclude that  $B_k(x(t, j)) = 0$  for all  $t \in [t'_1, t'_2]$  and the contradiction follows.  $\square$

**Remark 13** *As in Theorem 4, when the barrier function candidate  $B$  is scalar and locally Lipschitz, the statement of Proposition 1 holds true if we replace (29) therein by*

$$\max_{\zeta \in \partial_C B(x)} \langle \zeta, \eta \rangle \leq \rho(B(x)) \quad \forall x \in (U(K) \setminus K) \cap C \text{ and} \\ \forall \eta \in F(x) \cap T_C(x). \quad (30)$$

### 3.5 From Pre-Invariance to Invariance

In the following statement, we show when a forward pre-invariant set  $K \subset C \cup D$  is forward invariant. The proof can be found in [30].

**Proposition 2** *A forward pre-invariant set  $K \in C \cup D$  is forward invariant if the solutions cannot escape in finite time inside  $K \cap C$  and, for any initial condition in the set  $(K \cap \partial C) \setminus D$ , a nontrivial flow exists.*

**Remark 14** *One can guarantee that the solutions do not have a finite escape time inside the set  $K \cap C$  when, for example, the set  $K \cap C$  is compact or when the flow map  $F$  is globally bounded in  $K \cap C$ .*

**Example 8 (Thermostat)** *Using Proposition 2, we extend the conclusions in Example 2 and show that the set  $K$  introduced therein is forward invariant. Indeed, we already showed that the set  $K$  is forward pre-invariant. Furthermore, since it is compact, then there is not a possibility of a finite-time escape inside  $K \cap C$ . Finally, we note that  $(K \cap \partial C) \setminus D = (K \cap C) \setminus D = \{0\} \times [z_{min}, z_{max}] \cup \{1\} \times [z_{min}, z_{max}]$  and, by explicitly solving the flow dynamics on  $[0, t_1]$ , for some  $t_1 > 0$ , we conclude that, when  $x_o \in \{0\} \times [z_{min}, z_{max}]$ , there exists a nontrivial flow given by  $q(t, 0) = 0$ ,  $z(t, 0) = (z(0, 0) - z_o)e^{-t} + z_o$ . Similarly, when  $x_o \in \{1\} \times [z_{min}, z_{max}]$ , there exists a nontrivial flow given by  $q(t, 0) = 1$ ,  $z(t, 0) = (z(0, 0) - z_o - z_\Delta)e^{-t} + z_o + z_\Delta$ .*

In the following result, we propose a qualitative condition implying the existence of a nontrivial solution starting from each element in the set  $(K \cap \partial C) \setminus D$ .

**Proposition 3** *A forward pre-invariant set  $K \in C \cup D$  is forward invariant if the solutions cannot escape in finite time inside the set  $K \cap C$  and there exists a neighborhood  $U(x_o)$  such that*

$$F(x) \cap T_{K \cap C}(x) \neq \emptyset \quad \forall x \in U(x_o) \cap (K \cap \partial C) \text{ and} \\ \forall x_o \in (K \cap \partial C) \setminus D. \quad (31)$$

**PROOF.** To conclude the proof in this case, we propose to show that

$$T_{K \cap C}(x) \cap F(x) \neq \emptyset \quad \forall x \in U(x_o) \cap \partial(K \cap C) \text{ and} \\ \forall x_o \in (K \cap \partial C) \setminus D. \quad (32)$$

Indeed, using (32) and [30, Proposition 8] with the set  $K$  therein replaced by  $K \cap C$ , we conclude the existence of a nontrivial flow starting from each  $x_o \in (K \cap \partial C) \setminus D$ . Hence, the forward invariance of the set  $K$  follows using Proposition 2. Now, in order to show (32), we distinguish two complementary situations. First, when  $x \in \partial(K \cap C) \cap \partial C = K \cap \partial C$ , in this case, (32) follows

from (31). Second, when  $x \in \partial(K \cap C) \cap \text{int}(C)$ , in this case, since the set  $K$  is forward pre-invariant and since there exist always a nontrivial solution flowing from each  $x \in \text{int}(C)$  under the standing assumptions, we conclude the existence of a nontrivial solution flowing from  $x$  and remaining in  $K$  for a nontrivial interval of time. Hence, using [30, Proposition 7], (32) follows also when  $x \in \partial(K \cap C) \cap \text{int}(C)$ , which completes the proof.  $\square$

**Example 9 (Bouncing ball)** *For the system in Example 3, we will show forward invariance of the set  $K$  using Proposition 3. Indeed, the set  $K$  is compact; hence, the solutions starting from  $K$  cannot escape in finite time. Furthermore,  $(K \cap \partial C) \setminus D = \{0\} \times (0, 1]$  and, for any  $x_o \in (K \cap \partial C) \setminus D$ , we can take  $U(x_o) \cap (K \cap \partial C) = \{0\} \times [x_o/2, 1]$ . Next, for any  $x \in U(x_o) \cap (K \cap \partial C)$  with  $U(x_o)$  small enough, we notice that  $F(x)$  is transversal to  $\partial C$  and at the same time  $\langle \nabla B(x), F(x) \rangle \leq 0$ ; hence, using [30, Corollary 3], we conclude that  $F(x) \in D_C(x) \cap T_{K_e}(x)$ , which implies, using [30, Lemma 5], that  $F(x) \in T_{C \cap K}(x)$  and (31) follows.*

#### 4 Sufficient Conditions for Pre-Contractivity and Contractivity Using Barrier Functions

A pre-contractive set is a forward pre-invariant set such that whenever a solution starts from its boundary, it immediately leaves it and evolves towards its interior. The study of contractive sets is very important since several techniques to derive Lyapunov functions are based on the construction of contractive sets, the resulting Lyapunov functions are known as *set-induced Lyapunov functions*, see [8, 10–12].

##### 4.1 Definitions

A definition of contractivity for the particular sets named  $C$ -sets is proposed in [8] using the *Minkowski* functional, also named *gauge* function, for both differential and difference equations, see Definitions 3.3 and 3.4 in [8], respectively. The latter approach can be extended for general hybrid inclusions to define pre-contractivity and contractivity for  $C$ -sets. Indeed, we start recalling that a set  $K \subset C \cup D$  is said to be a  $C$ -set if it is compact, convex and includes the origin in its interior. Moreover, the corresponding Minkowski functional  $\Psi_K : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$  is given by

$$\Psi_K(x) := \inf \{ \mu \geq 0 : x \in \mu K \}. \quad (33)$$

**Definition 5 (Pre-contractivity for  $C$ -sets)** *A  $C$ -set  $K \subset C \cup D$  is said to be pre-contractive if*

$$\limsup_{h \rightarrow 0^+} \frac{\Psi_K(x + \eta h) - 1}{h} < 0 \quad \forall x \in \partial K \cap C, \quad \forall \eta \in F(x) \cap T_C(x), \quad (34)$$

$$\Psi_K(\eta) < 1 \quad \forall x \in D \cap K, \quad \forall \eta \in G(x). \quad (35)$$

**Definition 6 (Contractivity for  $C$ -sets)** *A  $C$ -set  $K \subset C \cup D$  is said to be contractive if it is pre-contractive and, in addition, starting from each element in the set  $(K \cap \partial C) \setminus D$ , a nontrivial solution exists.*

The following lemma establishes important consequences of the contractivity properties, in Definitions 5 and 6, on the behavior of the system's solutions. Based on these consequences, we will define pre-contractivity and contractivity for general closed sets that are not necessarily  $C$ -sets. The proof is in [30].

**Lemma 1** *If a  $C$ -set  $K \subset (C \cup D)$  is pre-contractive (respectively, contractive) according to Definition 5 (respectively, Definition 6), then it is forward pre-invariant (respectively, forward invariant) and, for any  $x_o \in \partial K$  and any nontrivial solution  $x$  starting from  $x_o$ , there exists  $T > 0$  and  $J \in \mathbb{N}^*$  such that  $x(t, j) \in \text{int}(K)$  for all  $(t, j) \in \text{dom } x \cap ([0, T] \times \{0\}) \cup (\{0\} \times \{0, 1, \dots, J\})$ ,  $(t, j) \neq (0, 0)$ .*

For general closed sets, we cannot use the Minkowski functional to define the contractivity notions since  $K$  may not be convex. Consequently, a trajectory-based definition, in the case of differential inclusions, is proposed in [9] under the name of *strict invariance*. In this section, we propose definitions of contractivity and pre-contractivity for hybrid systems based on the behavior of the solutions after reaching the boundary of the considered set. The aim of the proposed definitions is to preserve the properties established in Lemma 1. Furthermore, sufficient conditions in terms of barrier function candidates defining the (closed) set  $K$  are proposed.

**Definition 7 (Pre-contractivity for general sets)**

*A closed set  $K \subset C \cup D$  is said to be pre-contractive if it is forward pre-invariant and for every  $x_o \in \partial K$  and every nontrivial solution  $x$  starting from  $x_o$ , there exists  $T > 0$  and  $J \in \mathbb{N}^*$  such that  $x(t, j) \in \text{int}(K)$  for all  $(t, j) \in \text{dom } x \cap ([0, T] \times \{0\}) \cup (\{0\} \times \{0, 1, \dots, J\})$ ,  $(t, j) \neq (0, 0)$ .*

**Definition 8 (Contractivity for general sets)** *A closed set  $K \subset C \cup D$  is said to be contractive if it is pre-contractive and forward invariant.*

**Remark 15** *It is useful to notice that, in the particular case of differential inclusions, the pre-contractivity of a closed set  $K \subset \mathbb{R}^n$  reduces to the nonexistence of a (non-hybrid) solution  $t \mapsto x(t)$  starting from any  $x_o \in \partial K$  such that  $x([0, T]) \subset \mathbb{R}^n \setminus \text{int}(K)$  for some  $T > 0$ .*

##### 4.2 Pre-Contractivity

Next, we propose to characterize contractivity notions using barrier functions defining general closed sets<sup>4</sup>.

<sup>4</sup> The characterization for the case of  $C$ -sets is treated in the report version [30].

Our approach is mainly based on [30, Lemma 4], which characterizes the Dubovitsky-Miliutin cone  $D_{\text{int}(K)}$  at the boundary of the considered closed set in terms of the barrier function candidate defining the set. Furthermore, the latter fact is combined with [30, Theorem 6] in order to conclude contractivity.

**Theorem 5** *Given a hybrid system  $\mathcal{H} = (C, F, D, G)$  and a  $C^1$  barrier function candidate  $B$  defining the set  $K$  in (6). The set  $K$  is pre-contractive if, for each  $i \in \{1, 2, \dots, m\}$ ,*

$$\langle \nabla B_i(x), \eta \rangle < 0 \quad \forall x \in M_i \cap C \\ \forall \eta \in F(x) \cap T_C(x), \quad (36)$$

$$F(x) \cap T_{\partial C \cap \partial K}(x) = \emptyset \quad \forall x \in \partial K \cap \partial C, \quad (37)$$

$$B(\eta) < 0 \quad \forall x \in K \cap D \quad \forall \eta \in G(x), \quad (38)$$

$$G(x) \subset C \cup D \quad \forall x \in K \cap D, \quad (39)$$

$$G(x) \subset \text{int}(C \cup D) \quad \forall x \in \partial K \cap D. \quad (40)$$

**PROOF.** We consider, without loss of generality, the hybrid extension of  $\mathcal{H}$  denoted by  $\mathcal{H}_e := (C_e, F_e, D, G)$  where  $C_e := U(C) \subset \mathbb{R}^n$  and  $F_e : C_e \rightrightarrows \mathbb{R}^n$  is any extension of  $F$  to  $C_e$  that preserves the standing assumptions. Also, we recall that  $K_e = \{x \in \mathbb{R}^n : B(x) \leq 0\}$  and  $K_{ei} := \{x \in \mathbb{R}^n : B_i(x) \leq 0\}$ ; hence,  $K_e = \bigcap_{i=1}^m K_{ei}$ . We start noticing that for each  $x_o \in \partial K_e \cap C$  there exists  $I_{x_o} \subset \{1, 2, \dots, m\}$  such that  $x_o \in M_i$  if and only if  $i \in I_{x_o}$ . That is, we have  $x_o \in \bigcap_{i \in I_{x_o}} (\partial K_{ei} \cap C)$  and, also,  $x_o \in \bigcap_{i \in I_{x_o}} (\partial K_{ei} \cap \text{int}(C_e))$ . Next, using [30, Lemma 4] under (36), it follows that  $F_e(x_o) \cap T_C(x_o) \subset D_{\text{int}(K_{ei})}(x_o)$  for all  $i \in I_{x_o}$ . Moreover, when  $i \notin I_{x_o}$ , we also have  $F_e(x_o) \cap T_C(x_o) \subset D_{\text{int}(K_{ei})}(x_o)$  since in this case  $D_{\text{int}(K_{ei})}(x_o) = \mathbb{R}^n$ . Hence, we conclude that  $F_e(x_o) \cap T_C(x_o) \subset D_{\text{int}(K_e)}(x_o)$  since  $\bigcap_{i=1}^m D_{\text{int}(K_{ei})}(x_o) = D_{\text{int}(\bigcap_{i=1}^m K_{ei})}(x_o)$ .

As a second step, we show that  $F(x_o) \cap T_{C \setminus \text{int}(K_e)}(x_o) = \emptyset$ . Indeed, for each  $x \in \partial K_e \cap C$  and  $\eta \in F(x)$ , either  $\eta \notin T_C(x)$ , hence  $\eta \notin T_{C \setminus (\text{int}(K_e) \cap C)}(x) = T_{C \setminus \text{int}(K_e)}(x)$ , or  $\eta \in D_{\text{int}(K_e)}(x)$  which implies that  $\eta \notin T_{\mathbb{R}^n \setminus \text{int}(K_e)}(x)$  thus  $\eta \notin T_{C \setminus \text{int}(K_e)}(x)$ . Now, using [30, Proposition 7], we conclude the non existence of any solution  $x$  to  $\mathcal{H}_e$  starting from  $x_o$  and flowing in  $C \setminus \text{int}(K_e)$  along a non-trivial time interval. Hence, if a nontrivial flow  $x$  exists starting from  $x_o$ , then  $x(t, 0) \in C_e \setminus (C \setminus \text{int}(K_e))$  for all  $(t, 0) \in \text{dom } x \cap ([0, T] \times \{0\}) \setminus (0, 0)$  and for some  $T > 0$ . That is, for each solution  $x$  to  $\mathcal{H}_e$  flowing from  $x_o$ , either there exists  $T > 0$  such that  $x((0, T], 0) \subset \text{int}(K_e)$  or, there exists  $T > 0$  such that  $x((0, T], 0) \subset C_e \setminus C$ . Particularly, when  $x_o \in \partial K_e \cap \text{int}(C) = \partial K \cap \text{int}(C)$ , a non-trivial flow always exists, hence,  $x((0, T], 0) \in \text{int}(K)$

for some  $T > 0$ . Furthermore, when  $x_o \in \partial C \cap \partial K$ , using [30, Proposition 7] under (37), we conclude that  $x((0, T], 0) \cap (\partial K \cap \partial C) = \emptyset$  for some  $T > 0$ . In other words, the set  $\partial K \cap \partial C$  is not weakly forward invariant under (37). Hence, using the fact that  $\partial K \cap C = (\partial K_e \cap C) \cup (\partial K \cap \partial C)$ , it follows that each solution  $x$  flowing from  $x_o \in \partial K \cap C$  there exists  $T > 0$  such that either  $x((0, T], 0) \subset \text{int}(K_e) \setminus (\partial C \cap \partial K) \cap C$ , or  $x((0, T], 0) \subset C_e \setminus C$ . Going back to  $\mathcal{H}$ , the latter scenario is excluded; hence, for any solution flowing from  $x_o \in \partial K \cap C$ , there exists  $T > 0$  such that, if  $\text{dom } x \setminus (\mathbb{R}_{\geq 0}, 0) \neq \emptyset$ , then  $x(t, 0) \subset \text{int}(K)$  for all  $(t, 0) \in \text{dom } x \cap ([0, T] \times \{0\}) \setminus (0, 0)$ . On the other hand, under (38), (39), and the continuity of  $B$ , we conclude that for any  $x_o \in K \cap D$ ,  $G(x_o) \subset \text{int}(K_e) \cap (C \cup D) \subset K$ ; hence, all the possible jumps from  $K \cap D$  maintain the solution in the set  $K$ . Furthermore, using (40), we conclude that, for each  $x_o \in \partial K \cap D$  and after any possible jump, the solutions jump to the interior of the set  $K$ . Hence, if a jump is possible when starting from  $x_o \in \partial K \cap D$ , then  $x(t, j) \in \text{int}(K)$  for all  $(t, j) \in \text{dom } x \cap (\{0\} \times \{0, 1\}) \setminus (0, 0) \neq \emptyset$ ; which completes the proof.  $\square$

**Example 10** *Consider the hybrid system  $\mathcal{H}$  given by*

$$F(x) := \begin{bmatrix} -(x_2 + 1) \\ -2(x_2 + 1) + x_1 \end{bmatrix} \quad \forall x \in C,$$

$$C := \{x \in \mathbb{R}^2 : x_2 \in [0, 1], |x_1| \leq \sqrt{3}\},$$

$$G(x) := \frac{1}{\sqrt{3}} \begin{bmatrix} x_1 \\ \sqrt{3} \\ 2 \end{bmatrix} \quad \forall x \in D,$$

$$D := \{x \in \mathbb{R}^2 : x_2 = 0, |x_1| \leq \sqrt{3}\}.$$

*We will show that the set*

$$K := \{x \in \mathbb{R}^2 : x_1^2 + (x_2 + 1)^2 \leq 4, x_2 \geq 0\},$$

*which is not a  $C$ -set, is pre-contractive. Indeed, the set  $K$  can be defined using the  $C^1$  barrier function candidate  $B(x) := [x_1^2 + (x_2 + 1)^2 - 4 \quad -x_2]^\top$ . To verify (38)-(40), we note that, for each  $x \in K \cap D$ ,  $G(x) = [\frac{x_1}{\sqrt{3}} \quad \frac{1}{2}]^\top \in \text{int}(C \cup D)$  since  $(1/\sqrt{3})x_1 \leq 1$  for all  $x \in K \cap D = \{x \in \mathbb{R}^2 : x_2 = 0, |x_1| \leq \sqrt{3}\}$ ; hence (39)-(40) hold. Moreover,  $B(G(x)) = [x_1^2/3 - 7/4 \quad -1/2]^\top < 0$  since  $|x_1| \leq \sqrt{3}$  for all  $x \in K \cap D$ ; hence, (38) also follows. Furthermore,  $\langle \nabla B_1(x), F(x) \rangle = -4(x_2 + 1)^2 < 0$  for all  $x \in C \cap M_1 = \{x \in \mathbb{R}^2 : x_2 \geq 0, |[x_1 \quad x_2 + 1]| = 2\}$ . Next, for every  $x \in M_2 \cap C = \{x \in \mathbb{R}^2 : x_2 = 0, |x_1| \leq \sqrt{3}\}$ ,  $F(x) = [-1 \quad -2 + x_1]^\top \notin T_C(x)$  since  $F_2(x) = -2 + x_1 < 0$  for all  $x \in M_2 \cap C$ ; hence, (36) is satisfied. To show that (37) holds, we note that  $\partial K \cap \partial C = \{[0 \quad 1]^\top\} \cup M_2$ . Furthermore, when  $x = [0 \quad 1]^\top$ ,  $F(x) = [-2 \quad -4]^\top \notin T_{\partial K \cap \partial C}(x) = 0$ , and, when  $x \in M_2$ , we have already shown that  $F(x) \notin T_C(x)$ . The two latter facts imply that  $F(x) \notin T_{\partial C \cap \partial K}(x)$  for*

all  $x \in \partial K \cap \partial C$ . Hence, pre-contractivity of the set  $K$  follows according to Theorem 5.

The following result extends Theorem 5 when the barrier function candidate defining the closed set  $K$  is only locally Lipschitz. The proof is in [30].

**Theorem 6** *Given a hybrid system  $\mathcal{H} = (C, F, D, G)$  and a scalar locally Lipschitz barrier function candidate  $B$  defining the set  $K$  in (6). The set  $K$  is pre-contractive if (37)-(40) hold and*

$$\max_{\zeta \in \partial_C B(x)} \langle \zeta, \eta \rangle < 0 \quad \forall x \in K_e \cap C, \quad \forall \eta \in F(x) \cap T_C(x). \quad (41)$$

### 4.3 From Pre-Contractivity to Contractivity

In the sequel, we complement the sufficient conditions in Theorem 5 and Theorem 6 to conclude contractivity rather than only pre-contractivity.

**Proposition 4** *A pre-contractive closed set  $K \subset C \cup D$  is contractive provided that its solutions from  $K$  do not escape in finite time inside  $K \cap C$  and, starting from each initial condition in  $(K \cap \partial C) \setminus D$ , a nontrivial flow exists.*

In the following statement, the existence of a nontrivial flow starting from  $(K \cap \partial C) \setminus D$  is guaranteed provided that a qualitative tangentiality condition holds.

**Proposition 5** *Consider a  $C^1$  barrier function candidate defining the set  $K$  as in (6). The set  $K$  is contractive if conditions (36)-(40) hold and there exists a neighborhood  $U(x_o)$  such that*

$$F(x) \cap T_C(x) \neq \emptyset \quad \forall x \in U(x_o) \cap K \cap \partial C, \quad (42) \\ \forall x_o \in (K \cap \partial C) \setminus D.$$

**PROOF.** To conclude the proof, we show that

$$T_{K \cap C}(x) \cap F(x) \neq \emptyset \quad \forall x \in U(x_o) \cap \partial(K \cap C) \quad (43) \\ \forall x_o \in (K \cap \partial C) \setminus D.$$

Using [30, Proposition 8], (43) implies the existence of a nontrivial flow starting from each  $x_o \in (K \cap \partial C) \setminus D$ . Under the stated assumptions, the latter fact allows to conclude contractivity of the set  $K$  using Proposition 4. Now, to prove (43), we distinguish three complementary situations. First, when  $x \in \partial(K \cap C) \cap \partial C \cap \text{int}(K_e) = \partial C \cap \text{int}(K_e)$ . In this case, (43) follows from (42) since in this case  $T_C(x) = T_{K \cap C}(x)$ . Second, when  $x \in \partial(K \cap C) \cap \partial C \cap \partial K_e = \partial C \cap \partial K_e$ , in this case, we use (42) to conclude the existence of  $\eta \in F(x)$  such that  $\eta \in T_C(x)$ . Furthermore, we use (36) under [30, Lemma

4], to conclude that  $\eta \in \text{int}(T_{K_e}(x)) = D_{\text{int}(K_e)}(x)$ , which implies, using [30, Lemma 5], that  $\eta \in T_{K \cap C}(x)$ . Finally, consider  $x \in \partial(K \cap C) \cap \text{int}(C)$ . In this case, using Theorem 5, the set  $K$  is pre-contractive, hence, forward pre-invariant. Combining the latter fact to the existence of a nontrivial flow starting from each  $x \in \text{int}(C)$  under our standing assumptions, the existence of a nontrivial flow starting from  $x$  and remaining in  $K$  follows. Thus, under [30, Proposition 7], (43) follows when  $x \in \partial(K \cap C) \cap \text{int}(C)$ .  $\square$

**Example 11** *Using Proposition 5, We will show that the closed set  $K$  introduced in Example 10 is contractive. To this end, we need to show that there is not a possibility of finite-time escape inside  $K$ . This is the case since the set  $K$  is compact. Finally, to show that (42) holds for all  $x \in (K \cap \partial C) \setminus D = \{(0, 1)\}$ , we note that we can choose  $U(x) \cap K \cap \partial C = \{x := [0 \ 1]^T\}$  on which  $F(x) = [-2 \ -4]^T \in T_C(x)$  since  $F_2(x) = -4 < 0$ .*

## 5 Conclusion

The first part of this paper proposed sufficient conditions for forward invariance of closed sets for hybrid systems modeled as hybrid inclusions. The considered closed sets are defined using barrier function candidates and the proposed sufficient conditions in terms of the latter barrier functions are infinitesimal inequalities; namely, not involving any knowledge about the system's solutions, guaranteeing that the set of points on which all the components of the barrier function candidate are nonpositive is forward invariant. Studying forward invariance in the general context of hybrid inclusions offered many technical challenges that have not been handled in the existing literature, to the best of our knowledge. Those challenges are mainly due to the continuous-time evolution of the hybrid inclusion being not necessarily defined on an open set. Hence, elements around the intersection between the zero-level sets of the barrier candidate and the boundary of the set where the continuous-time evolution is defined needed a particular treatment. Finally, we proposed relaxed flow conditions compared to the existing literature by using the notion of uniqueness functions. In the following, we recap the different sufficient conditions for forward pre-invariance proposed in this paper. Given a hybrid system  $\mathcal{H} = (C, F, D, G)$  and a barrier function candidate  $B$  defining the closed set  $K$  in (6). The set  $K$  is forward pre-invariant for  $\mathcal{H}$  if (12)-(13) hold and either

1. (27) holds, or
2.  $B \in C^1$  and (11) holds, or
3.  $B \in C^1$  and (29) holds, or
4.  $B$  is locally Lipschitz and (28) holds, or
5.  $B$  is locally Lipschitz and (30) holds, or
6.  $B \in C^1$ , Assumption 1, (20), and (16) hold, and either (a), (b), or (c) in Theorem 2 holds.

In the second part of the paper, following the same approach as in the first part, we proposed sufficient conditions for contractivity, which is a stronger property than forward invariance.

In the future, it would be interesting to analyze the necessity of the different sufficient conditions proposed in this paper or to propose new necessary and sufficient ones. Also, it would be interesting to analyze the notions considered in this paper in the presence of perturbations. Furthermore, this work constitutes an important step to analyze the mixed safety plus convergence problem in hybrid systems. Indeed, the latter problem is solved if we show forward invariance of the safety region plus contractivity of the reachable set from the safety region towards a given target. Investigating sufficient (infinitesimal) conditions to guarantee the latter mixed safety-convergence task in hybrid systems is part of our current research efforts.

## References

- [1] P. Wieland and F. Allgöwer. Constructive safety using control barrier functions. *IFAC Proceedings Volumes*, 40(12):462–467, 2007.
- [2] G. S. Ladde and V. Lakshmikantham. On flow-invariant sets. *Pacific Journal of Mathematics*, 51(1):215–220, 1974.
- [3] A. Taly and A. Tiwari. Deductive verification of continuous dynamical systems. In *Proceedings of the LIPICs-Leibniz International Proceedings in Informatics*, volume 4. Schloss Dagstuhl-Leibniz-Zentrum für Informatik, 2009.
- [4] S. Prajna. *Optimization-based methods for nonlinear and hybrid systems verification*. PhD thesis, California Institute of Technology, 2005.
- [5] R. G. Sanfelice, R. Goebel, and A. R. Teel. Invariance principles for hybrid systems with connections to detectability and asymptotic stability. *IEEE Transactions on Automatic Control*, 52(12):2282–2297, 2007.
- [6] R. G. Sanfelice and A. R. Teel. Asymptotic stability in hybrid systems via nested matrosov functions. *IEEE Transactions on Automatic Control*, 54(7):1569–1574, 2009.
- [7] G. S. Ladde and S. Leela. Analysis of invariant sets. *Annali di Matematica Pura ed Applicata*, 94(1):283–289, 1972.
- [8] F. Blanchini. Survey paper: Set invariance in control. *Automatica*, 35(11):1747–1767, 1999.
- [9] J. P. Aubin. *Viability Theory*. Birkhauser Boston Inc., Cambridge, MA, USA, 1991. ISBN 0-8176-3571-8.
- [10] M. Fiacchini, C. Prieur, and S. Tarbouriech. On the computation of set-induced control lyapunov functions for continuous-time systems. *SIAM Journal on Control and Optimization*, 53(3):1305–1327, 2015.
- [11] F. Blanchini. Ultimate boundedness control for uncertain discrete-time systems via set-induced lyapunov functions. *IEEE Transactions on Automatic Control*, 39(2):428–433, Feb 1994. ISSN 0018-9286. doi: 10.1109/9.272351.
- [12] P. Giesl and S. Hafstein. Review on computational methods for Lyapunov functions. *Discrete and Continuous Dynamical Systems-Series B*, 20(8):2291–2331, 2015.
- [13] M. Nagumo. Über die lage der integralkurven gewöhnlicher differentialgleichungen. *Proceedings of the Physico-Mathematical Society of Japan. 3rd Series*, 24:551–559, 1942.
- [14] F. H. Clarke, Y. S. Ledyaev, R. J. Stern, and P. R. Wolenski. *Nonsmooth Analysis and Control Theory*, volume 178. Springer Science & Business Media, 2008.
- [15] J. P. Aubin, L. Lygeros, M. Quincampoix, S. Sastry, and N. Seube. Impulse differential inclusions: A viability approach to hybrid systems. *IEEE Transactions on Automatic Control*, 47(1):2–20, 2002.
- [16] J. Chai and R. G. Sanfelice. Forward invariance of sets for hybrid dynamical systems (Part I). *IEEE Transactions on Automatic Control*, 64(6):2426–2441, 2018.
- [17] A. D. Ames, X. Xu, J. W. Grizzle, and P. Tabuada. Control barrier function based quadratic programs for safety critical systems. *IEEE Transactions on Automatic Control*, 62(8):3861–3876, 2017.
- [18] P. Glotfelter, J. Cortés, and M. Egerstedt. Nonsmooth barrier functions with applications to multi-robot systems. *IEEE control systems letters*, 1(2):310–315, 2017.
- [19] P. Glotfelter, I. Buckley, and M. Egerstedt. Hybrid nonsmooth barrier functions with applications to provably safe and composable collision avoidance for robotic systems. *IEEE Robotics and Automation Letters*, 4(2):1303–1310, 2019.
- [20] S. Prajna, A. Jadbabaie, and G. J. Pappas. A framework for worst-case and stochastic safety verification using barrier certificates. *IEEE Transactions on Automatic Control*, 52(8):1415–1428, 2007.
- [21] X. Xu, J. W. Grizzle, P. Tabuada, and A. D. Ames. Correctness guarantees for the composition of lane keeping and adaptive cruise control. *IEEE Transactions on Automation Science and Engineering*, 15(3):1216–1229, 2018.
- [22] Q. Nguyen and K. Sreenath. Safety-critical control for dynamical bipedal walking with precise footstep placement. *IFAC-PapersOnLine*, 48(27):147–154, 2015.
- [23] L. Dai, T. Gan, B. Xia, and N. Zhan. Barrier certificates revisited. *Journal of Symbolic Computation*, 80:62–86, 2017. ISSN 0747-7171. SI: Program Verification.
- [24] H. Kong, F. He, X. Song, W. N. N. Hung, and M. Gu. Exponential-condition-based barrier certificate generation for safety verification of hybrid




systems. In *Proceedings of the Computer Aided Verification*, pages 242–257, Berlin, Heidelberg, 2013. Springer Berlin Heidelberg.

- [25] M. Z. Romdlony and B. Jayawardhana. Stabilization with guaranteed safety using control lyapunov-barrier function. *Automatica*, 66:39–47, 2016.
- [26] F. H. Clarke. *Optimization and Nonsmooth Analysis*, volume 5. 1990.
- [27] M. Maghenem and R. G. Sanfelice. Barrier function certificates for forward invariance in hybrid inclusions. In *Proceedings of the 56th IEEE Conference on Decision and Control (CDC)*, pages 759–764, 2018.
- [28] M. Maghenem and R. G. Sanfelice. Multiple barrier function certificates for forward invariance in hybrid inclusions. In *Proceedings of the 2019 American Control Conference (ACC)*, pages 2346–2351. IEEE, 2019.
- [29] M. Maghenem and R. G. Sanfelice. Barrier function certificates for forward invariance in hybrid inclusions. In *Proceedings of the 56th IEEE Conference on Decision and Control (CDC)*, pages 759–764, 2018.
- [30] M. Maghenem and R. G. Sanfelice. Sufficient conditions for forward invariance and contractivity in hybrid inclusions using barrier functions. *arXiv preprint:1908.03980*, 2019.
- [31] R. Goebel, R. G. Sanfelice, and A. R. Teel. Hybrid dynamical systems. *IEEE Control Systems magazine*, 29(2):28–93, 2009.
- [32] J. P. Aubin and H. Frankowska. *Set-valued Analysis*. Springer Science & Business Media, 2009.
- [33] R. Goebel, R. G. Sanfelice, and A. R. Teel. *Hybrid Dynamical Systems: Modeling, stability, and robustness*. Princeton University Press, 2012.
- [34] S. Boyd and L. Vandenberghe. *Convex optimization*. Cambridge university press, 2004.
- [35] H. G. Tanner, A. Jadbabaie, and G. J. Pappas. Stable flocking of mobile agents, Part I: Fixed topology. In *Proceedings of the 42nd Conference on Decision and Control*, volume 2, pages 2010–2015. IEEE, 2003.
- [36] A. G. Wills and W. P. Heath. Barrier function based model predictive control. *Automatica*, 40(8):1415–1422, 2004.
- [37] M. Jankovic. Robust control barrier functions for constrained stabilization of nonlinear systems. *Automatica*, 96:359–367, 2018.
- [38] J. M. Bony. Principe of the maximum, inégalité de harnack et unicité du problème de cauchy pour les operateurs elliptiques dégénérés. *Ann. Inst. Fourier (Grenoble)*, 19(1):277–304, 1969.
- [39] H. Brezis. On a characterization of flow-invariant sets. *Communications on Pure and Applied Mathematics*, 23(2):261–263, 1970.
- [40] R. M. Redheffer. The theorems of bony and brezis on flow-invariant sets. *The American Mathematical*

*Monthly*, 79(7):740–747, 1972.

- [41] R. P. Agarwal and V. Lakshmikantham. *Uniqueness and nonuniqueness criteria for ordinary differential equations*. World Scientific Publishing Company, 1993.

## Biography



**Mohamed Maghenem** received his Control-Engineer degree from the Polytechnical School of Algiers, Algeria, in 2013, his M.S. and Ph.D. degrees in Control from the University of Paris-Saclay, France, in 2014 and 2017, respectively. He is currently a Postdoctoral Fellow at the Electrical and

Computer Engineering Department at the University of California at Santa Cruz. His research interests include dynamical systems theory (stability, safety, reachability, robustness, and synchronization), control systems theory (adaptive, time-varying, linear, non-linear, hybrid, robust, etc.) with applications to power systems, mechanical systems, and cyber-physical systems.



**Ricardo G. Sanfelice** received the B.S. degree in Electronics Engineering from the Universidad de Mar del Plata, Buenos Aires, Argentina, in 2001, and the M.S. and Ph.D. degrees in Electrical and Computer Engineering from the University of California, Santa Barbara, CA, USA, in 2004 and 2007, respectively. In 2007 and

2008, he held postdoctoral positions at the Laboratory for Information and Decision Systems at the Massachusetts Institute of Technology and at the Centre Automatique et Systèmes at the Ecole de Mines de Paris. In 2009, he joined the faculty of the Department of Aerospace and Mechanical Engineering at the University of Arizona, Tucson, AZ, USA, where he was an Assistant Professor. In 2014, he joined the University of California, Santa Cruz, CA, USA, where he is currently Professor in the Department of Electrical and Computer Engineering. Prof. Sanfelice is the recipient of the 2013 SIAM Control and Systems Theory Prize, the National Science Foundation CAREER award, the Air Force Young Investigator Research Award, the 2010 IEEE Control Systems Magazine Outstanding Paper Award, and the 2020 Test-of-Time Award from the Hybrid Systems: Computation and Control Conference. His research interests are in modeling, stability, robust control, observer design, and simulation of nonlinear and hybrid systems with applications to power systems, aerospace, and biology.